

Par III®

HERBICIDE

Par III the most trusted name in golf course and turf management, is a broad-spectrum herbicide specially formulated to control hard-to-kill annual and perennial weeds in turfgrass. Par III is ideal for golf courses, parks, lawns, sports fields and other turf applications where weeds cause aesthetic and durability problems. Used in a twice-yearly maintenance program, Par III keeps turfgrass free of tough problem weeds such as dandelion, poison ivy, clovers and ground ivy.

ACTIVE INGREDIENT: mecoprop (d-isomer 100 g/L); 2,4-D (isomer specific 190 g/L); dicamba 18 g/L (all present as dimethylamines).

CROPS: Turf.

RATE: For turfgrass (not containing bentgrass) use 5.5 L of Par III in 300 L water per hectare. (55 ml in 3 L water per 100 m²).

BENTGRASS TURF: On turf containing closely mowed bentgrass (putting and bowling greens, etc.), use a maximum rate of 30 ml in 20 L of water per 100 m² to prevent injury - preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

SPOT TREATMENTS: For spot treatments, use 200 ml in 10 L water and wet foliage thoroughly using a backpack sprayer.

WATER VOLUME: Sufficient water for adequate coverage using low pressure (200-275 kPa).

TANK MIXES: None.

TIMING: Par III can be applied to established turf at any time during the growing season when weeds are young and actively growing. Apply in good growing conditions before crop canopy shields the weeds. Do not apply to newly established turf until after second to third mowings. If used prior to seeding as a pre-seed weed control, grass can be sown 1-2 weeks after application.

Spring and early fall provide the best opportunity for maximum weed control. Avoid application during long, excessively dry or hot periods unless adequate irrigation is available.



ALWAYS READ THE LABEL BEFORE USE

CAUTION: Do not apply more than two applications per year not including spot treatments. Do not apply by air. Avoid spray drift to any desirable vegetation not listed on label. Avoid contamination of ponds, streams and other water sources. Do not spray in high winds.

RAINFAST: Do not irrigate within 24 hours or spray within 24 hours of rainfall.

PREHARVEST INTERVAL: For best weed control, avoid mowing for several days before and after treatment.

RE-ENTRY TIME: Do not allow people (other than the applicator), or pets on treated area during application. Do not re-enter until thoroughly dried.

PACKAGE SIZE: Par III is available in a range of package sizes: 12 X 1L; 2 X 10 L and 115 L.

RESIDUAL CONTROL: No soil residual activity on weeds. Two applications per season are all that is needed under good turf/lawn management.

ADDITIONAL: Use of flat fan nozzle recommended. Ensure sprayer is clean before use. Use a tank cleaner such as All Clear™ to remove unwanted residues in the sprayer tank.

SURFACTANT NEEDED: None.

STORAGE: Store tightly sealed containers away from seeds, feeds, fertilizers, plants and foodstuffs. Protect from freezing. Product exposed to subfreezing temperatures should be warmed to 5_ C and mixed thoroughly before using.

HERBICIDE GROUP: Group 4.

WEEDS CONTROLLED

Bedstraw	Heal all
Black medick (yellow clover)	Knotweed (ironweed)
Buttercup	Orange hawkweed (devil's paint brush)
Chickweed	Plantain
Clover	Poison ivy
Dandelion	Ragweed
English daisy	Shepherd's purse
Ground ivy (creeping Charlie)	



www.uap.ca

WEST: 1-800-561-5444 Ontario & Maritimes: 1-800-265-5444 Quebec: 1-800-361-9369