

# Pounce®

## INSECTICIDE

Pounce is a broad spectrum insecticide with low mammalian toxicity. Pounce has good residual activity to help minimize the need for repeat applications. It controls insects through a combination of stomach and contact activity, which improves control effectiveness.

**TARGET PESTS:** Many common insect pests of fruit, vegetable, greenhouse and ornamental crops. Please see label for complete listing.

Armyworms  
Caterpillar type larvae  
Cutworms (various species)  
European corn borer  
Fleabeetle  
Mushroom flies  
Potato leafhoppers  
Whitefly

**CROPS:** Wide range of crops, (see label for specifics).

**Field crops** including cereals, canola, corn, flax, lentils, peas, potatoes, and sunflowers.

**Vegetables** including cole crops, sweet corn, cucumber (greenhouse) and peppers.

**Fruit** including apple, blueberries, pears, peaches, grapes, nectarines and plums.

**Other crops and uses** including mushroom, tobacco, farm buildings, ornamental and greenhouse use.

**RATE:** 45 ml-1.0 L/ha depending on crop (see label).

**WATER VOLUME:** Dependent on crop type, please consult label. Apply sufficient water volume to achieve good coverage as there is no systemic or fumigant effect. Use higher rates for heavy infestations, adult insects and dense foliage.

**GROUP:** Group 3; synthetic pyrethroid insecticide.

**TIMING:** Application timing depends on target insect. Various use patterns include pre-plant soil applied; post-plant soil applied; broadcast spray when insects or insect damage first appears. See label for specific recommendations regarding insect, application timing and days to harvest restrictions.

Consult with your local crop advisor or Provincial Crop Protection Guides to determine economic threshold levels when controlling insects.

**CAUTION:** Do not use when outdoor temperature exceeds 25° C as effectiveness could be reduced.

**RAINFAST:** Once dried, with optimum of 24 hours to allow insects to feed.

**PREHARVEST INTERVAL:** 1-21 days depending on crop and use. Cover crops or crops treated with Pounce should not be used as green feed for animals.



## QUICK SHEET

