

Products



Ridomil Gold® 480SL Fungicide

Uses

CROPS

Potatoes; Refer to the label for a complete list of crops.

FOR MANAGEMENT OF

For suppression of the following disease in

Potatoes: In-furrow: Pink rot (Phytophtho...

[View more use info](#)

Application Information

Potatoes: In-furrow: 4 mL/100 m row Foliar: 208 mL/ha (80 mL/ac) One jug of R...

Water Volume

Potatoes: Foliar - Apply in a minimum of 30 L of water/ha (12 L/ac or 3 gal/ac).

[View more application info](#)

Tank Mixes



[View more tank-mixes info](#)

Technical Information

CHEMISTRY GROUP
Group 4 fungicide

ACTIVE INGREDIENTS
Metalaxyl-M and S-isomer

REI: **12 hours**

[View more technical info](#)

Label and MSDS

Label

[Download](#)

MSDS

[Download](#)

[View more labels](#)

About this Product

Help mitigate financial loss in storage

Incorporate Ridomil Gold® 480SL fungicide into your pink rot management strategy. Consider tank-mixing with Allegro® for increased activity on late blight, and protection against white mould. Growers should follow through with a post-harvest application of a registered product targeted at pink rot and tuber blight going into storage.

Packaging

Case: 4 x 3.78 L jugs

Uses

- Potatoes

Refer to the label for a complete list of crops.

For management of

For suppression of the following disease in Potatoes:

In-furrow:

- Pink rot (*Phytophthora erythroseptica*)*

Foliar:

- Pink rot (*Phytophthora erythroseptica*)*
- Late blight (*Phytophthora infestans*)*
- Pythium leak

*There is known Ridomil Gold 480SL resistance to certain strains of late blight and pink rot. Ridomil Gold 480SL should always be tank-mixed with another product that controls late blight when targeting that disease.

Refer to the label for a complete list of crops and diseases.

Application Information

Use Rates

Potatoes:

- In-furrow: 4 mL/100 m row
- Foliar: 208 mL/ha (80 mL/ac)

One jug of Ridomil Gold 480SL treats 45 acres as a foliar application OR 22.5 acres as an in-furrow application.

Refer to the label for a complete listing of crops, diseases and rates.

Water Volume

Potatoes:

Foliar - Apply in a minimum of 30 L of water/ha (12 L/ac or 3 gal/ac).

Application Method

Drench,Ground,Other

Application Information

Potatoes:

- Apply in-furrow or in two foliar applications.
- When applying to foliage, use in a tank-mix with another late blight fungicide (we recommend Allegro®).
- Make a first application at tuber initiation, then a second application 14 days later.

Refer to the label for more application details, including those for other crops.

Use Restrictions

Potatoes:

- PHI: In-furrow: 80 days
- PHI: Foliar: 14 days
- REI: 12 hours

Refer to the label for a complete list of use restrictions for all crops.

Tank Mixes

For Potatoes:

In-furrow

- Actara
- Minecto Duo
- Quadris

Foliar

- Allegro
- Bravo ZN
- Quadris Top
- Revus

Refer to the label for a complete list of tank mix partners and the application details for each.

Technical Information

Chemistry Group

Group 4 fungicide

Mode of Action

Ridomil Gold 480SL contains the active ingredient Metalaxyl-M and S-isomers. Ridomil Gold 480SL is readily absorbed by the roots and provides protection against attack from these diseases under varied weather conditions. The metalaxyl-m in Ridomil Gold is an RNA polymerase inhibitor.

Labels & MSDS

Label

[Download](#)

Description For the control of certain diseases caused by the Oomycete class of fungi in Apples (non-bearing), Field and Greenhouse Cucumbers,

Ginseng, Hops, Raspberries, Blueberries, Snap Beans, Strawberries, Tobacco and Potatoes.

Active ingredient Metalaxyl-M and S-isomer

Approved provinces AB, BC, MB, NB, NF, NS, ON, PE, QC, SK

Approved crops Apples (nonbearing), Blueberries, Cucumber, Ginseng, Hops, Potatoes, Radish, Raspberries, Spinach, Strawberries, Tobacco

MSDS

[Download](#)

Disclaimer – ALWAYS read and follow label directions. Please use this information above only as a guideline and not an exact description.

© 2015 Syngenta