

GROUP	6	HERBICIDE
-------	---	-----------

BASAGRAN® FORTE HERBICIDE

LIQUID

For selective post-emergence broadleaf weed control in soybeans, dry beans, peas, corn (grain, silage, sweet and seed), flax (excluding low linolenic acid varieties), sorghum and millet (grain and forage), fababeans, and established red and alsike clover for seed production

COMMERCIAL (AGRICULTURAL)

GUARANTEE: Bentazon480 g/L

Warning, contains the allergen soy

REGISTRATION NO. 22006

PEST CONTROL PRODUCTS ACT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**



WARNING

POISON

**DANGER – CORROSIVE TO EYES
POTENTIAL SKIN SENSITIZER**

NET CONTENTS: 10 L

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING
SHAKE WELL BEFORE USE
KEEP OUT OF REACH OF CHILDREN**

BASF Canada Inc.
100 Milverton Drive
5th Floor
Mississauga, Ontario
L5R 4H1
1-877-371-2273

GENERAL INFORMATION

BASAGRAN FORTE Herbicide is a liquid herbicide for selective post-emergence control of many broadleaf weeds and yellow nutsedge in soybeans, dry beans, peas, corn (grain, silage, sweet and seed), flax (excluding low linolenic acid varieties), fababeans, forage and grain sorghum, forage and grain millet, and established red and alsike clover for seed production. For enhanced broad spectrum weed control in soybeans only, **BASAGRAN FORTE Herbicide** can be tank mixed with a **nitrogen source** such as **28% UAN**, **PINNACLE®**, **PURSUIT®** or **BLAZER®**, depending upon target weeds.

BASAGRAN FORTE Herbicide does not control grasses.

BASAGRAN FORTE Herbicide is a herbicide with mainly contact action. Uptake into the plant occurs primarily through the leaves. Thorough coverage of foliage is important for consistent weed control. Failure to penetrate crop or weed leaf canopies with the spray will result in incomplete control of small weeds growing underneath.

Cool weather conditions or drought will delay herbicidal activity and if prolonged, may result in poor weed control. Use of **BASAGRAN FORTE Herbicide** in hot, humid weather may result in temporary leaf yellowing, leaf flecking, bronzing or burning particularly on peas and some varieties of dry common beans. The crop usually outgrows this condition within 10 days (see Restrictions and Limitations).

SPRAYING INSTRUCTIONS

Ground Application

Use sprayers equipped with standard flat fan pesticide nozzles with the recommended spray volume, pressure and additives. Tilt spray nozzles 45 degrees forward to ensure better coverage.

SPRAY VOLUME	SPRAY PRESSURE
Minimum 100 L/ha. Use larger water volumes (up to 300 L/ha) for weeds at the upper limit of their recommended stage for treatment.	Minimum 275 kPa. Use higher pressure (use up to 425 kPa) for weeds at the upper limit of their recommended stage for treatment.

NOTE: Use a minimum of 200 L/ha if crop canopy or heavy weed population interferes with thorough spray coverage, or under cool temperature.

BASAGRAN FORTE HERBICIDE BANDING TECHNIQUES

1. Spray a minimum of 25 cm wide band.
2. Minimize the amount of dust striking target weeds to ensure adequate coverage and penetration.
3. **DO NOT** use cultivation equipment when spraying.
4. Adjust the **BASAGRAN FORTE Herbicide** rate to proportion of the total area to be sprayed.

SPRAY DRIFT MANAGEMENT FOR GROUND APPLICATION

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

Buffer Zones

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, rangelands, riparian areas and shrublands).

Method of Application	Crop	Buffer zone (metres) required for the protection of:
		Terrestrial Habitat
Field sprayer ¹	Sorghum and millet (grain and forage), red and alsike clover	1
	Soybeans, corn, dry beans, fababeans, peas, flax	2

¹ For field sprayer application, buffer zones can be reduced with the use of drift-reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

DIRECTIONS FOR USE

Timing of Application

Although the timing of application should be primarily in relation to the stage of weed growth, the crop must be in a tolerant stage as shown in Table 1. The information on usual stage of crop for optimum weed control may also serve as a guide to spray timing. (Refer to Table 1 - Crop Application Timing.)

Early treatment of weeds with **BASAGRAN FORTE Herbicide** is important to maximize crop yield potential through elimination of early weed competition and permits better coverage of the weeds, thereby increasing the performance of **BASAGRAN FORTE Herbicide**.

Apply **BASAGRAN FORTE Herbicide** at the correct rate when broadleaf weeds are small and actively growing and before the weeds reach the maximum size recommended for treatment as listed in Tables 2, 3 and 4. **BASAGRAN FORTE Herbicide** should be applied when the main weed of concern is in the suggested growth stage for treatment.

Table 5 details the tank mix options available for soybeans and the relative maximum weed leaf stages or weed height in centimetres for effective control. For detailed information on specific tank mixes, refer to Tank Mix Sections for each tank mix partner.

TABLE 1 - BASAGRAN FORTE HERBICIDE CROP APPLICATION TIMING

CROP	TOLERANCE STAGE	USUAL STAGE OF CROP FOR OPTIMUM WEED CONTROL
Beans (dry common) ¹ – <i>Phaseolus vulgaris</i> only	Tolerant after 1st trifoliolate leaf has fully expanded	1 to 3 trifoliolate leaves
Corn (grain, silage, sweet and seed)	Tolerant at any growth stage	1 to 5 leaf stage
Fababeans	Tolerant after 2 or 3 leaf stage or crop is 10 cm high	Soon after 3 leaf stage
Flax (excluding low linolenic acid varieties)	Tolerant when crop is 5 cm or higher	Soon after crop reaches 5 cm
Sorghum and millet (grain and forage) ²		3 to 6 leaf stage before closure of crop canopy
Peas (field and processing)	Tolerant after 3 pairs of leaves (or 3 nodes) are present but prior to crop flowering	Soon after 3 pairs of leaves form
Soybeans	Tolerant at any growth stage	Unifoliolate to two expanded trifoliolate leaves usually 18 to 28 days after planting
Forage legumes (established, for seed production) ^{2,3} o Red and alsike clover	Tolerant between 7.5 and 25 cm high	After crop is 7.5 cm and before crop canopy closes.

¹ Dry common bean varieties may vary in their tolerance to herbicides, including **BASAGRAN FORTE Herbicide**. Since not all dry common bean varieties have been tested for tolerance to **BASAGRAN FORTE Herbicide**, first use of **BASAGRAN FORTE Herbicide** should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific varieties of dry common beans to **BASAGRAN FORTE Herbicide**.

² Apply one application/year only.

³ Crop injury may occur under hot, humid conditions. Speed of recovery will be influenced by growing conditions and weed control.

TABLE 2 - BASAGRAN FORTE HERBICIDE WEED APPLICATION TIMING

TREATMENT	WEED SPECIES CONTROLLED	MAXIMUM LEAF STAGE
BASAGRAN FORTE Herbicide at 2.25 L/ha	Bird rape*	6-leaf
	Buttercup	6-leaf
	Chickweed (Common)	1-3 weeks after emergence
	Cleavers	1-3 whorls
	Cocklebur	10-leaf
	Corn spurry	10 cm
	Flower-of-an-hour	10-leaf
	Galinsoga (Hairy)	6-leaf
	Groundsel (Common)*	10 cm
	Jimsonweed	10-leaf
	Lady's-thumb (Smartweed)	10-leaf
	Lamb's-quarters*	8-leaf
	Low cudweed	6-leaf
	Mustard (Wild)	10-leaf
	Nightshade (Hairy)	6-leaf
	Pigweed (Redroot)* (suppression only)	4-leaf
	Purslane	6-leaf
	Radish (Wild)	6-leaf
	Ragweed (Common)*	6-leaf
	Ragweed (Giant)	4-leaf
	Shepherd's-purse	6-leaf
Stinkweed	6-leaf	
Thistle (Russian) (suppression only)	4-leaf	
Velvetleaf**	6-leaf	

TABLE 3 - BASAGRAN FORTE HERBICIDE WEED APPLICATION TIMING

TREATMENT	WEED SPECIES CONTROLLED	MAXIMUM LEAF STAGE
BASAGRAN FORTE Herbicide at 1.75 L/ha	Bird rape*	6-leaf
	Cocklebur	6-leaf
	Flower-of-an-hour	6-leaf
	Lady's-thumb (Smartweed)	6-leaf
	Mustard (Wild)	6-leaf
	Shepherd's-purse	6-leaf
	Stinkweed	6-leaf
	Velvetleaf**	4-leaf
Perennial Weeds: Repeat application 7 - 15 days after first treatment (if necessary)		
BASAGRAN FORTE Herbicide at 1.75 L/ha	Bindweed (Field)*** (suppression only)	6 cm
	Thistle (Canada)	20 cm
	Yellow nutsedge	20 cm

* Triazine resistant strains of these weeds are controlled by **BASAGRAN FORTE Herbicide**.

** **BASAGRAN FORTE Herbicide** will defoliate velvetleaf 4 leaf or larger, but regrowth may occur.

*** Treat field bindweed before it is dark green and has begun to trail.

TABLE 4 – BASAGRAN FORTE HERBICIDE WEED APPLICATION TIMING FOR FIELD PEAS GROWN IN ALBERTA

TREATMENT	WEED SPECIES CONTROLLED	MAXIMUM LEAF STAGE
BASAGRAN FORTE Herbicide at 1.25 L/ha*	Spiny annual sow-thistle (suppression only)	4-leaf
BASAGRAN FORTE Herbicide at 1 L/ha*	Volunteer canola (including all herbicide tolerant varieties)	4-leaf

*Apply one application/year only.

TABLE 5 - TANK MIX WEED APPLICATION TIMING (SOYBEANS ONLY)

The following table details the tank mix options available and the relative maximum **weed leaf stage or weed height in centimetres** for application. For detailed information on specific tank mixes, refer to Tank Mix Sections for each tank mix partner.

WEEDS (Leaf stages)	BASAGRAN FORTE Herbicide + PURSUIT + N Source***	BASAGRAN FORTE Herbicide + PURSUIT + N Source***	BASAGRAN FORTE Herbicide + BLAZER	BASAGRAN FORTE Herbicide + BLAZER	BASAGRAN FORTE Herbicide + PINNACLE	BASAGRAN FORTE Herbicide + PINNACLE
RATE	1.25 L/ha + 0.312 L/ha + 2 L/ha	1.75 L/ha + 0.312 L/ha + 2 L/ha	1.75 L/ha + 0.63 L/ha	1.25 L/ha + 1.25 L/ha	1.75 L/ha + 5.5 or 8.0 g/ha	2.25 L/ha + 5.5 or 8.0 g/ha
Bindweed (Field) (Suppression)**		6 cm			6 cm	6 cm
Bird rape*		6 leaf	6 leaf		6 leaf	6 leaf
Buttercup						6 leaf
Chickweed (Common)						1-3 weeks after emergence
Cleavers						1-3 whorls
Cocklebur	4 leaf	6 leaf	6 leaf		6 leaf	10 leaf
Corn spurry						10 cm
Flower-of-an-hour		6 leaf	6 leaf		6 leaf	10 leaf
Galinsoga (Hairy)						6 leaf
Groundsel (Common)						10 cm
Jimsonweed						10 leaf
Lady's thumb (Smartweed)		6 leaf	8 leaf		6 leaf	10 leaf
Lamb's-quarters	6 leaf	6 leaf	6 leaf	6 leaf	8 leaf	8 leaf
Low cudweed						6 leaf
Mustard (Wild)	3 leaf	6 leaf	8 leaf		6 leaf	10 leaf
Nightshade (Eastern Black)	4 leaf	4 leaf	4 leaf			

WEEDS (Leaf stages)	BASAGRAN FORTE Herbicide + PURSUIT + N Source***	BASAGRAN FORTE Herbicide + PURSUIT + N Source***	BASAGRAN FORTE Herbicide + BLAZER	BASAGRAN FORTE Herbicide + BLAZER	BASAGRAN FORTE Herbicide + PINNACLE	BASAGRAN FORTE Herbicide + PINNACLE
RATE	1.25 L/ha + 0.312 L/ha + 2 L/ha	1.75 L/ha + 0.312 L/ha +2 L/ha	1.75 L/ha + 0.63 L/ha	1.25 L/ha + 1.25 L/ha	1.75 L/ha + 5.5 or 8.0 g/ha	2.25 L/ha + 5.5 or 8.0 g/ha
Pigweed (Redroot)*	12 leaf	12 leaf	8 leaf	8 leaf	6 leaf	6 leaf
Purslane						6 leaf
Radish (Wild)						6 leaf
Ragweed (Common)*	4 leaf	6 leaf	8 leaf	8 leaf		6 leaf
Ragweed (Giant)						4 leaf
Shepherd's-purse		6 leaf	6 leaf		6 leaf	6 leaf
Stinkweed		6 leaf	6 leaf			6 leaf
Thistle (Canada)**		20 cm			20 cm	20 cm
Thistle (Russian) (Suppression only)						4 leaf
Velvetleaf	8 leaf	8 leaf	4 leaf		4 leaf	6 leaf
Yellow nutsedge**		20 cm			20 cm	20 cm
Annual grass Including:						
Barnyard grass	6 leaf	6 leaf				
Foxtail (Green and yellow)	4 leaf	4 leaf				

* Triazine resistant strains of these weeds are controlled by **BASAGRAN FORTE Herbicide**.

** May require an additional application of the 1.75 L/ha rate of **BASAGRAN FORTE Herbicide** only for control.

*** Nitrogen sources include: 10-34-0, 28-0-0 or 32-0-0.

TANK MIXTURES

ALL TANK MIXES ARE REGISTERED FOR USE IN SOYBEANS ONLY. USE OF THESE TANK MIXES IN CROPS OTHER THAN SOYBEANS MAY RESULT IN SIGNIFICANT INJURY OR LOSS OF CROP.

BASAGRAN FORTE Herbicide + NITROGEN SOURCE

Benefit (over BASAGRAN FORTE Herbicide alone):

Improved control of velvetleaf and lamb's-quarters.

Rate:

Add 6 L/ha ammonium sulphate or 10 L/ha 28% urea ammonium nitrate (UAN) to either the 2.25 L/ha or the 1.75 L/ha rate of **BASAGRAN FORTE Herbicide**.

Notes:

The addition of ammonium sulphate or 28% urea ammonium nitrate to **BASAGRAN FORTE Herbicide** will result in more consistent control of velvetleaf and lamb's-quarters, especially under cool conditions, heavy weed pressures or at the maximum growth stages indicated in the **BASAGRAN FORTE Herbicide** Weed Application Timing (Tables 2 and 3).

The addition of either form of nitrogen source to **BASAGRAN FORTE Herbicide** may cause slight leaf burn, but the new growth is normal and crop vigour is not reduced. A high quality source of 28% UAN or ammonium sulphate should be used to avoid solids or contaminants that may interfere with spray application (clogging nozzle tips) or reduced product performance. Do not use brass or aluminum nozzles when spraying **BASAGRAN FORTE Herbicide** plus a nitrogen source.

BASAGRAN FORTE HERBICIDE + PURSUIT + NITROGEN SOURCE (SOYBEANS ONLY)

Benefit (over BASAGRAN FORTE Herbicide alone):

Two modes of action providing postemergent contact and soil activity. Control of nightshade (Eastern black), redroot pigweed, annual grasses and improved control of lamb's-quarters.

Rate:

**BASAGRAN FORTE Herbicide 1.25 L/ha or
1.75 L/ha + PURSUIT 0.312 L/ha + Nitrogen Source 2 L/ha**

Add 0.312 L/ha of **PURSUIT** to 1.25 L/ha or 1.75 L/ha rate of **BASAGRAN FORTE Herbicide** + 2 L/ha of nitrogen source for control of nightshade (Eastern black), velvetleaf, common

ragweed, redroot pigweed, barnyard grass, green and yellow foxtail, and improved control of lamb's-quarters.

When weeds are under stress due to environmental conditions, use the high rate of **BASAGRAN FORTE Herbicide** at 1.75 L/ha + **PURSUIT** at 0.312 L/ha + nitrogen source at 2 L/ha.

WEED SPECIES CONTROLLED	MAXIMUM LEAF STAGE
Lamb's quarters	6
Nightshade (Eastern black)	4
Pigweed (redroot)	12
Ragweed (common)	4
	Basagran Forte Herbicide 1.25 L/ha
	*Basagran Forte Herbicide 1.75 L/ha
Velvetleaf	8
Annual Grasses	
Barnyard grass	6
Green foxtail	4
Yellow foxtail	4

For additional weeds controlled by the tank mix of **BASAGRAN FORTE Herbicide** and **PURSUIT**, consult Table 4 for maximum leaf stages or weed heights.

* Use 1.75 L/ha rate of **BASAGRAN FORTE Herbicide** for control of additional weeds and later stages, as detailed in Table 4, including cocklebur, lady's thumb, and wild mustard.

Use 1.75 L/ha rate of **BASAGRAN FORTE Herbicide** for perennial weed control, including yellow nutsedge, Canada thistle and field bindweed (suppression only). A second application of **BASAGRAN FORTE Herbicide** alone at 1.75 L/ha may be required 7-15 days after initial treatment.

Notes:

Slight initial crop stunting may result when tank mixing with **PURSUIT** but will not result in yield reduction or delay in harvest.

When tank mixing **BASAGRAN FORTE Herbicide** with **PURSUIT**, a pre-harvest interval of 100 days should be maintained.

Always refer to **PURSUIT** label for rotational crop restrictions, first aid, precautions, warning and disposal of herbicide.

BASAGRAN FORTE HERBICIDE + BLAZER

Benefit (over BASAGRAN FORTE Herbicide alone):

Improved control of ragweed, redroot pigweed and nightshade.

Rate:

Add 1.25 L/ha **BLAZER** to 1.25 L/ha rate of **BASAGRAN FORTE Herbicide**.

BLAZER at 1.25 L/ha plus **BASAGRAN FORTE Herbicide** at 1.25 L/ha will control the weeds listed in Table 4 up to the specified leaf stage. Use this treatment where common ragweed and/or redroot pigweed are the dominant weed species present.

OR

Add 0.63 L/ha **BLAZER** to 1.75 L/ha rate of **BASAGRAN FORTE Herbicide**.

BLAZER at 0.63 L/ha plus **BASAGRAN FORTE Herbicide** at 1.75 L/ha will control the weeds listed in Table 4 up to the specified leaf stage. Use this treatment where lamb's-quarters and weeds other than common ragweed and redroot pigweed are the dominant weed species present.

Note:

Do not use ASSIST Oil Concentrate or other adjuvants with the BLAZER plus BASAGRAN FORTE Herbicide tank mixture.

BASAGRAN FORTE HERBICIDE + PINNACLE (SOYBEANS ONLY)

Benefit (over BASAGRAN FORTE Herbicide alone):

Control of redroot pigweed and improved control of lamb's-quarters.

Rate:

BASAGRAN FORTE Herbicide 2.25 L/ha + PINNACLE 5.5 or 8.0 g/ha
BASAGRAN FORTE Herbicide 1.75 L/ha + PINNACLE 5.5 or 8.0 g/ha

Add **PINNACLE** at 5.5 or 8.0 g/ha to either the 2.25 L/ha or the 1.75 L/ha rate of **BASAGRAN FORTE Herbicide** for improved control of certain broadleaf weeds, including lamb's-quarters and redroot pigweed as well as all the weeds listed in the corresponding "**BASAGRAN FORTE Herbicide Weed Application Timing Tables 2 and 3**" for **BASAGRAN FORTE Herbicide**

alone. The following table details the leaf stages where control of redroot pigweed and lamb's-quarters can be expected.

All weeds controlled by **BASAGRAN FORTE Herbicide** plus the following:

WEEDS SPECIES CONTROLLED	MAXIMUM LEAF STAGE
Redroot pigweed	6-leaf
+Lamb's-quarters	8-leaf

+Use 8.0 g/ha rate of **PINNACLE** for advanced leaf staging.

- * For control of common ragweed, use the 2.25 L/ha rate of **BASAGRAN FORTE Herbicide**.
- * For additional weeds controlled by the tank mix of **BASAGRAN FORTE Herbicide** and **PINNACLE**, consult Table 4 for maximum weed leaf stages or weed heights.

Notes:

The addition of **PINNACLE** to **BASAGRAN FORTE Herbicide** provides excellent control of redroot pigweed and lamb's-quarters. However, even though redroot pigweed growth is rapidly stopped, typical symptoms of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions.

The combination of **BASAGRAN FORTE Herbicide** and **PINNACLE** does not provide residual weed control. Weeds that emerge after application will not be controlled.

Do not add nitrogen source or any other additive when tank mixing **BASAGRAN FORTE Herbicide** with **PINNACLE**.

Always refer to the **PINNACLE** label for use, precautions, sprayer cleanup, first aid, storage and disposal of **PINNACLE**.

MIXING

1. Clean spray tank by following sprayer cleanup recommendations on the previous product label and fill half full with clean water. Start agitation or by-pass system.
2. When required add correct amount of Nitrogen source. **Note: Do not add nitrogen source when tank mixing BASAGRAN FORTE Herbicide with PINNACLE.**
3. If required, add the correct amount of **PINNACLE** or **PURSUIT**. **Note:** For use on soybeans only. Continuous agitation is required to keep **PINNACLE** or **PURSUIT** in suspension.

Note: When using the **BASAGRAN FORTE Herbicide + PINNACLE** tank mix, prepare a **PINNACLE/water slurry** in a separate container with clean water before adding to spray tank on repeat tank loads.

4. Add correct amount of **BASAGRAN FORTE Herbicide** and agitate 2 to 3 minutes.
5. Add remainder of water, agitate and spray.
6. If an oil film starts to build up in the tank, drain and clean tank with a strong detergent solution.
7. Thoroughly clean spray tank immediately after use. Refer to specific tank mix partner label for cleanup instructions.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **BASAGRAN FORTE Herbicide** is a Group 6 herbicide. Any weed population may contain or develop plants naturally resistant to **BASAGRAN FORTE Herbicide** and other Group 6 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **BASAGRAN FORTE Herbicide** or other Group 6 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact BASF at 1-877-371-2273.

RESTRICTIONS AND LIMITATIONS

1. **DO NOT** treat any crops not listed on this label.
2. **DO NOT** use **BASAGRAN FORTE Herbicide** with additives, pesticides or fertilizers not specifically recommended on this label.
3. **DO NOT** apply **BASAGRAN FORTE Herbicide** to any crops that have been subjected to stress from conditions such as hail damage, flooding, drought, hot, humid weather, widely fluctuating temperature conditions, prolonged cold weather or injury from prior herbicide applications, as crop injury may result.
4. Rainfall within 6 to 8 hours of application may reduce effectiveness of spray.
5. When **BASAGRAN FORTE Herbicide** is applied beyond recommended weed growth stages, limited to unsatisfactory weed control will result.
6. Cool weather conditions or drought will delay herbicide activity and if prolonged, may result in poor weed control.
7. Do not apply **BASAGRAN FORTE Herbicide** within 30 days of harvest for forage sorghum and forage millet, and within 100 days of harvest for grain sorghum and grain millet.
8. Do not apply **BASAGRAN FORTE Herbicide** within 36 days of harvest of red clover grown for seed production.
9. **DO NOT APPLY BY AIR.**

WARNING

1. **DO NOT** apply **BASAGRAN FORTE Herbicide** when weather conditions may cause spray drift from treated areas to adjacent crops.
2. Lentils, adzuki and mung beans, cucumbers, sugar beets and sunflowers can be injured by **BASAGRAN FORTE Herbicide**.

ENVIRONMENTAL HAZARDS

The use of this chemical may result in contamination of groundwater, particularly in areas where soils are permeable (e.g. sandy soil) or the water table is shallow.

DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), or to estuarine/marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

TOXIC to non-target terrestrial plants. Observe buffer zones specified under SPRAY DRIFT MANAGEMENT FOR GROUND APPLICATION.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. Harmful or fatal if swallowed or inhaled. Do not take internally.
3. Corrosive to the eye. Do not get in eyes.
4. Avoid inhalation of vapour, dust or spray mist and contact with eyes, skin or clothing. Potential skin sensitizer.
5. Wash thoroughly after handling and before eating, drinking or smoking.
6. Wear protective equipment and clothing, including: goggles or face shield, approved respirator, gloves (rubber, PVC, neoprene or nitrile), hat, long-sleeved shirt, trousers and rubber boots.
7. If clothing becomes contaminated, remove and wash separately from household laundry before re-use.
8. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
9. DO NOT enter or allow worker entry into treated areas for 12 hours following application.
10. Clean spray equipment thoroughly after use. When tank mixing **BASAGRAN FORTE Herbicide** with another herbicide, refer to respective label for sprayer cleanup.
11. **CAUTION: Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.**

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

FIRST AID INSTRUCTIONS

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

1. Store in original, tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross contamination.
5. Protect from freezing.

DISPOSAL

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For Designated Refill Containers

Container Refilling and Disposal: If this container is to be refilled, do not rinse with any material or introduce any pesticide other than **BASAGRAN FORTE Herbicide**. **DO NOT REUSE THIS CONTAINER FOR ANY OTHER PESTICIDE.** Reseal and return to any authorized BASF Refill Station. Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices. Check for leaks after filling and before transportation. Do not refill or transport damaged or leaking containers.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

ASSIST® and PURSUIT® are registered trademarks of BASF.

®All other products listed are registered trademarks of their respective companies.

GROUP

6

HERBICIDE

BASAGRAN® FORTE HERBICIDE

LIQUID

For selective post-emergence broadleaf weed control in soybeans, dry beans, peas, corn (grain, silage, sweet and seed), flax (excluding low linolenic acid varieties), sorghum and millet (grain and forage), fababeans, and established red and alsike clover for seed production

COMMERCIAL (AGRICULTURAL)

GUARANTEE: Bentazon480 g/L

Warning, contains the allergen soy

REGISTRATION NO. 22006

PEST CONTROL PRODUCTS ACT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**



WARNING

POISON

**DANGER – CORROSIVE TO EYES
POTENTIAL SKIN SENSITIZER**

NET CONTENTS: 10 L

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING
SHAKE WELL BEFORE USE
KEEP OUT OF REACH OF CHILDREN**

BASF Canada Inc.
100 Milverton Drive
5th Floor
Mississauga, Ontario
L5R 4H1
1-877-371-2273

BASAGRAN® FORTE is a registered trademark of BASF. ©BASF Canada Inc., 1992

PRECAUTIONS

- Wear protective equipment and clothing, including: goggles or face shield, approved respirator, gloves (rubber, PVC, neoprene or nitrile), hat, long-sleeved shirt, trousers and rubber boots.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- DO NOT enter or allow worker entry into treated areas for 12 hours following application.
- **CAUTION: Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.**
- **DO NOT APPLY BY AIR.**

Please refer to the booklet for further precautions.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

ENVIRONMENTAL HAZARDS

The use of this chemical may result in contamination of groundwater, particularly in areas where soils are permeable (e.g. sandy soil) or the water table is shallow.

DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), or to estuarine/marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

TOXIC to non-target terrestrial plants. Observe buffer zones specified under SPRAY DRIFT MANAGEMENT FOR GROUND APPLICATION.

FIRST AID INSTRUCTIONS

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

1. Store in original, tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross contamination.
5. Protect from freezing.

DISPOSAL

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

GROUP

2

HERBICIDE

Pursuit[®]
herbicide

FOR SOYBEANS

AGRICULTURAL

(NOT FOR SALE FOR USE IN PRAIRIE PROVINCES)

SOLUTION

GUARANTEE: imazethapyr ... 240 g/L

REGISTRATION No. 26287 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1.26 Litres

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT

1-800-454-2673

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

BASF Canada Inc.
100 Milverton Drive
5th Floor
Mississauga, Ontario
L5R 4H1
1-877-371-2273

PURSUIT is a registered trade-mark of BASF Agrochemical Products, B.V., used with permission by BASF Canada Inc.

©BASF Canada Inc., 2000

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.

DO NOT CONTAMINATE FOOD OR FEED PRODUCTS.

AVOID CONTACT WITH EYES, SKIN AND CLOTHING.

AVOID BREATHING VAPOUR OR SPRAY MIST.

May cause eye damage.

May cause skin irritation.

Use with adequate ventilation.

Wear dust and/or splash-proof goggles or face shield and chemically resistant gloves when mixing, loading and during application, clean-up and repair.

Wear long-sleeved shirt and long-legged pants when handling.

Do not eat, drink or smoke when using.

Wash hands and face before eating, drinking, smoking and using the toilet.

Take a shower immediately after work.

Store and wash all protective clothing separately from normal laundry.

Clean protective equipment (gloves, goggles, face shield) upon removal with soapy water.

Wash in detergent and hot water before reuse.

Wear freshly laundered clothes daily.

Do not enter or allow entry into treated areas during the restricted-entry interval of 12 hours.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

ENVIRONMENTAL HAZARDS

TOXIC to non-target terrestrial and aquatic plants. Observe buffer zones specified under DIRECTIONS FOR USE.

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Keep from freezing. DO NOT store below 0°C. If the product is exposed to temperatures below 0°C during shipment or storage, make sure the product has thawed completely, and shake the container vigorously.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE

CROP: SOYBEANS

GENERAL

PURSUIT is a selective herbicide that can be applied as a post-emergence treatment in soybeans. Absorption may occur through both the roots and foliage of susceptible weeds which stop growing and eventually die.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

Buffer Zones

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of Application	Crop	Buffer Zones (metres) Required for the Protection of:	
		Freshwater Habitat	Terrestrial Habitat
Field sprayer	Soybeans	1	1

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

ROTATIONAL CROPS

Only soybeans, kidney beans, cranberry, Dutch brown, black, yellow eye, white beans, adzuki beans, field corn, imazethapyr tolerant corn, imazethapyr tolerant canola, processing peas, spring barley, spring wheat and winter wheat may be planted the season following a PURSUIT herbicide application. Winter wheat may be grown the same season as a PURSUIT herbicide application to soybeans but not earlier than 100 days after the application. Conduct a field bioassay (a test strip grown to maturity) the year BEFORE growing any other crop.

RATE: Apply 312 - 420 mL of product (75 - 100 g active) in 100 to 400 litres of water per hectare.

EARLY POST-EMERGENCE APPLICATION

Early post-emergence applications of PURSUIT herbicide may be applied after the crop has emerged and before the weeds reach the 2 true leaf stage. **Addition of a non-ionic surfactant (AGRAL® 90 or AGSURF®) plus liquid fertilizer is essential for post-emergence application, improving uptake of the product by weeds resulting in improved herbicidal activity.** Application to a crop under stress should be avoided. Stunting of the crop may result following post-emergence application. This condition is the result of stem internode shortening but will not result in yield reduction.

BROADLEAF WEED AND GRASS CONTROL

PURSUIT herbicide, by early post-emergence application BEFORE the weeds reach the 2 true leaf stage, will control:

Broadleaf Weeds: redroot pigweed
velvetleaf*
wild mustard
cocklebur*/**
eastern black nightshade
ragweed**
wild buckwheat

Grasses: green foxtail
yellow foxtail
old witchgrass
barnyard grass**

In addition, PURSUIT herbicide provides partial control of and reduces competition from:

<u>Broadleaf Weeds:</u>	lamb's-quarters	<u>Grasses:</u>	proso millet large crabgrass
<u>Perennial Weed:</u>	yellow nutsedge		

* Some plants of cocklebur and velvetleaf which germinate deeper in the soil and emerge late may escape treatment.

** The higher label rate is recommended for fields with a history of heavy infestations of these weed species.

NOTE: For post-emergence applications a NONIONIC SURFACTANT (AGRAL 90 or AGSURF) at the rate of 0.25% (v/v) MUST BE ADDED to the spray solution (e.g. 2.5 L of surfactant per 1000 L of spray solution). Fertilizer solutions, 10-34-0, 28-0-0 or 32-0-0 MUST BE ADDED to the spray solution at the rate of 2 L/ha to provide quicker burndown of weeds.

EARLY POST-EMERGENCE TANKMIXING FOR BROAD SPECTRUM WEED CONTROL

For fields which contain heavy infestations of ragweed, velvetleaf, barnyard grass or weeds other than those listed under PURSUIT herbicide alone, tankmixing may be required to provide broad spectrum weed control.

PURSUIT herbicide may be tankmixed with either BASAGRAN® or BASAGRAN FORTE to provide a broader spectrum of weed control. Follow rates, recommendations, precautions and restrictions on the BASAGRAN and BASAGRAN FORTE labels.

NOTE: For PURSUIT herbicide plus BASAGRAN herbicide, a NON-IONIC SURFACTANT at the rate of 0.25% (v/v) and fertilizer solution (10-34-0, 28-0-0 or 32-0-0) at the rate of 2 L/ha MUST BE ADDED to the spray solution. For PURSUIT herbicide plus BASAGRAN FORTE herbicide only fertilizer solution must be added at the rate of 2 L/ha.

MIXING INSTRUCTIONS

Fill the tank one-half to three-quarters full with water and start agitation. Add the required amount of 28-0-0, 10-34-0, or 32-0-0 to the spray solution. Using a calibrated measuring device, add the required amount of PURSUIT herbicide while the solution remains agitating. Using a separate calibrated measuring device, add the required amount of BASAGRAN or BASAGRAN FORTE herbicide and make sure it is thoroughly mixed. Continue agitation while adding the required amount of non-ionic surfactant (for PURSUIT herbicide + BASAGRAN herbicide ONLY). Continue agitation while filling the remainder of the tank with water.

LATE POST-EMERGENCE APPLICATION

PURSUIT herbicide may be applied late post-emergence for control of certain weed species up to the growth stages indicated below. To minimize weed competition with the crop, application should be made as early as possible after weed emergence. Avoid applications when weeds and/or crop are under growth stress.

Addition of a non-ionic surfactant and liquid fertilizer is critical for post-emergence application, see instructions for tankmixing. Stunting of the crop may result following post-emergence application. This condition is the result of stem internode shortening but will not result in yield reduction.

BROADLEAF WEED AND GRASS CONTROL

PURSUIT herbicide by late post-emergence application will control:

<u>GRASS Weeds</u>	<u>Weed Species</u>	<u>Maximum Growth Stage</u>
	green foxtail	4 leaf
	yellow foxtail	4 leaf
	barnyard grass	6 leaf
<u>BROADLEAF Weeds</u>	<u>Weed Species</u>	<u>Maximum Growth Stage</u>
	redroot pigweed	12 leaf
	velvetleaf	8 leaf

LATE POST-EMERGENCE TANKMIXING FOR BROAD SPECTRUM WEED CONTROL

For broadleaf weeds other than those listed for PURSUIT herbicide alone, tankmix with BASAGRAN or BASAGRAN FORTE herbicide. Follow rate, recommendations, precautions and restrictions on the BASAGRAN or BASAGRAN FORTE herbicide label. Tankmix product and rate will depend on the specific weeds to be controlled.

* Use the high rate of (100 g ae/ha) 168 mL product/acre when PURSUIT herbicide is used alone. Use the (75 g ae/ha) 126 mL when tankmixed with BASAGRAN or BASAGRAN FORTE herbicide.

NOTE: For PURSUIT herbicide and PURSUIT herbicide plus BASAGRAN herbicide, a NON-IONIC SURFACTANT at the rate of 0.25% (v/v) and fertilizer solution (10-34-0, 28-0-0 or 6-24-6) at the rate of 2 L/ha MUST BE ADDED to the spray solution. For PURSUIT herbicide plus BASAGRAN FORTE herbicide only fertilizer solution must be added at the rate of 2 L/ha.

MIXING INSTRUCTIONS: LATE POST-EMERGENCE APPLICATIONS

Fill the spray tank one-half to three-quarters full with water and start agitation. Add the required amount of 28-0-0, 10-34-0, or 6-24-6 to the spray solution. Using a calibrated measuring device, add the required amount of PURSUIT herbicide while the solution remains agitating. Using a separate calibrated measuring device, add the required amount of BASAGRAN or BASAGRAN FORTE herbicide and make sure it is thoroughly mixed. Continue agitation while adding the required amount of non-ionic surfactant (for PURSUIT herbicide + BASAGRAN herbicide ONLY). Continue agitation while filling the remainder of the tank with water.

WARNINGS

ONLY soybeans, kidney beans, cranberry, Dutch brown, black, yellow eye, white beans, adzuki beans, field corn, imazethapyr tolerant corn, imazethapyr tolerant canola, processing peas, spring barley, spring wheat and winter wheat may be planted the season following a PURSUIT herbicide application. Winter wheat should not be planted within 100 days of the last PURSUIT herbicide application. Conduct a field bioassay (a test strip grown to maturity) the year BEFORE growing any other crop.

DO NOT overapply PURSUIT herbicide. Over application may result in injury particularly if the crop is under stress.

An interval of 100 days should follow the last PURSUIT herbicide application before soybean harvest.

CAUTION: Do not graze treated crops or cut for hay; sufficient data are not available to support such use.

DO NOT let spray drift contaminate crops other than those being sprayed.

ONLY ONE (1) application of PURSUIT herbicide may be made during the season.

In cases of crop failure, only soybeans, kidney beans, cranberry, Dutch brown, black, yellow eye, white beans, adzuki beans, imazethapyr tolerant canola, imazethapyr tolerant corn, processing peas and winter wheat (100 days after application) may be re-planted in the year of application. Soil preparation for re-planting should be no deeper than 10 cm.

DO NOT apply PURSUIT herbicide when crop is under stress conditions because crop injury may result.

PURSUIT herbicide may cause stunting of soybeans. This condition is the result of stem internode shortening and does not cause yield reductions if PURSUIT herbicide has been used at label rates and following label recommendations.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, PURSUIT is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to PURSUIT and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of PURSUIT or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact BASF at 1-877-371-2273.

BASAGRAN is a registered trade-mark of BASF SE, used with permission by BASF Canada Inc.

®Unless specified otherwise, products listed are registered trademarks of their respective companies.

GROUP

2

HERBICIDE

Pursuit[®]
herbicide

FOR SOYBEANS

AGRICULTURAL

(NOT FOR SALE FOR USE IN PRAIRIE PROVINCES)

SOLUTION

GUARANTEE: imazethapyr ... 240 g/L

REGISTRATION NO. 26287 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1.26 Litres

IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT

1-800-454-2673

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

BASF Canada Inc.
100 Milverton Drive
5th Floor
Mississauga, Ontario
L5R 4H1
1-877-371-2273

PURSUIT is a registered trade-mark of BASF Agrochemical Products, B.V., used with permission by BASF Canada Inc.

©BASF Canada Inc., 2000

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.

DO NOT CONTAMINATE FOOD OR FEED PRODUCTS.

AVOID CONTACT WITH EYES, SKIN AND CLOTHING.

AVOID BREATHING VAPOUR OR SPRAY MIST.

May cause eye damage.

May cause skin irritation.

Use with adequate ventilation.

Wear dust and/or splash-proof goggles or face shield and chemically resistant gloves when mixing, loading and during application, clean-up and repair.

Wear long-sleeved shirt and long-legged pants when handling.

Do not eat, drink or smoke when using.

Wash hands and face before eating, drinking, smoking and using the toilet.

Take a shower immediately after work.

Store and wash all protective clothing separately from normal laundry.

Clean protective equipment (gloves, goggles, face shield) upon removal with soapy water.

Wash in detergent and hot water before reuse.

Wear freshly laundered clothes daily.

Do not enter or allow entry into treated areas during the restricted-entry interval of 12 hours.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

DO NOT APPLY BY AIR.

ENVIRONMENTAL HAZARDS

TOXIC to non-target terrestrial and aquatic plants. Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Keep from freezing. DO NOT store below 0°C. If the product is exposed to temperatures below 0°C during shipment or storage, make sure the product has thawed completely, and shake the container vigorously.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.