

Box Label (Base)

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

During all activities, wear long pants, long-sleeved shirt and work boots. In addition, chemical resistant gloves, protective eyewear and work boots must be worn during mixing, loading, cleanup and repair of spray equipment.

For aerial application, coveralls or a chemical resistant apron over normal work clothes, chemical resistant gloves, protective eyewear and work boots must be worn during mixing and loading operations.

DO NOT re-enter treated areas within 12 hours. If required, individuals may re-enter treated areas within 12 hours for short-term tasks not involving hand labour if at least 4 hours have passed since application and long pants, long-sleeved shirt, hat, protective eyewear and chemical resistant gloves are worn.

Do not inhale product or spray. This product is mildly irritating to eyes. Wash hands and exposed skin before eating, drinking or smoking. Thoroughly clean application equipment after use. Do not contaminate domestic or irrigation waters. Do not store in contact with foodstuffs.

FIRST AID:

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically. Medical personnel should contact Bayer CropScience's medical information services, toll-free 1-800-334-7577.

DISPOSAL:

1. Empty bag.
2. Follow provincial instructions for any required additional cleaning of the bag prior to its disposal.
3. Make the empty bag unsuitable for further use.
4. Dispose of the bag in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

Box Label (Base)

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Booklet Label

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

During all activities, wear long pants, long-sleeved shirt and work boots. In addition, chemical resistant gloves, protective eyewear and work boots must be worn during mixing, loading, cleanup and repair of spray equipment.

For aerial application, coveralls or a chemical resistant apron over normal work clothes, chemical resistant gloves, protective eyewear and work boots must be worn during mixing and loading operations.

DO NOT re-enter treated areas within 12 hours. If required, individuals may re-enter treated areas within 12 hours for short-term tasks not involving hand labour if at least 4 hours have passed since application and long pants, long-sleeved shirt, hat, protective eyewear and chemical resistant gloves are worn.

Do not inhale product or spray. This product is mildly irritating to eyes. Wash hands and exposed skin before eating, drinking or smoking. Thoroughly clean application equipment after use. Do not contaminate domestic or irrigation waters. Do not store in contact with foodstuffs.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca.

CAUTION: Do not graze the treated crops or use for feed or fodder; sufficient data are not available to support such use.

AIR APPLICATION

- Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.
- Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.
- Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Aerial Use Precautions

- Read and understand the entire label before opening this product. If you have questions, call Bayer CropScience Inc. at 1-888-283-6847 or obtain technical advice from the distributor or your provincial agricultural representative.
- Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.
- Do not apply to terrain where there is a potential for surface run-off to enter aquatic systems.

Booklet Label

- **Do not apply if rain is expected within 1 hour after spraying. If tank mixing, consult the tank mix partner label and respect the most restrictive interval.**

Aerial Application Operator Precautions

- Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.
- It is desirable that the pilot has communication capabilities at each treatment site at the time of application.
- The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.
- All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

FIRST AID:

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically. Medical personnel should contact Bayer CropScience's medical information services, toll-free 1-800-334-7577.

DISPOSAL:

1. Empty bag.
2. Follow provincial instructions for any required additional cleaning of the bag prior to its disposal.
3. Make the empty bag unsuitable for further use.
4. Dispose of the bag in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Booklet Label

DIRECTIONS FOR USE:

For application through field, aerial or orchard sprayers, put one half the required water volume in the tank, stir in Rovral[®] and add the remainder of the water. For use through knapsack or small sprayers, first stir the required amount of Rovral[®] with water in a bucket, add this suspension to the partly filled spray tank, then add the remainder of the water.

Ensure mixing is complete and agitate thoroughly before each application and continue to agitate during spraying.

A prepared spray mixture should be used as soon as possible in the same day, especially if the water used for mixing is alkaline (greater than pH of 7).

Rovral[®] should not be mixed with other pesticides, adjuvants or fertilizers except where stated.

FRUIT AND FIELD CROPS

APRICOT- CONTROL OF BROWN ROT/BLOSSOM BLIGHT

RATE: 1.5 kg/ha of Rovral[®].

TIMING: To protect the fruit use Rovral[®] in a cover spray program with application at 7-14 day intervals throughout the season. Use the short term interval under wet conditions close to harvest. For blossom blight apply at first bloom. Repeat at 4-5 day intervals if weather is wet. Apply by ground. Observe a 1 day PHI. To reduce the possibility of disease resistance to Rovral[®], alternation with other fungicides is recommended.

CHERRY - CONTROL OF BROWN ROT

RATE: 1.5-1.75 kg/ha of Rovral[®]. Use the higher rate on large or mature trees. For dilute sprays use 50 g/100 litres of water.

TIMING: Blossom Blight Stage: Spray at first bloom. If wet weather prevails, this spray should be repeated.

Brown Rot: To protect the fruit use Rovral[®] in a spray program with applications beginning at Shuck. Repeat at 7-14 day intervals throughout the season, including a treatment 1 day prior to harvest. To reduce the possibility of disease resistance to Rovral[®], the alternation with other fungicides is recommended.

GRAPE - CONTROL OF BOTRYTIS BUNCH ROT

RATE: 1.5 kg/ha Rovral[®] in at least 900 litres of water per hectare.

TIMING: Sprays may be applied beginning at mid to late bloom. Two sprays should be applied before complete bunch closure; these should be timed to achieve maximum protection of the fruit, i.e. just before anticipated periods of wet weather. To reduce the possibility of disease

Booklet Label

resistance to Rovral[®], no more than two applications of Rovral[®] should be made per season; other fungicides should be alternated with Rovral[®] in a bunch rot control program.

PEACH - CONTROL OF BROWN ROT

RATE: 1.5-1.75 kg/ha of Rovral[®]. Use the higher rate on large or mature trees. For dilute sprays, use 50 g/100 litres of water.

TIMING: Blossom Blight: Spray at first bloom to control blossom blight stage of brown rot. Repeat at 4-5 day intervals if the weather is wet.

Brown Rot: To protect the fruit use Rovral[®] in a cover spray program with application at 7-14 day intervals throughout the season including a treatment 1 day prior to harvest. Use the short interval under wet conditions close to harvest. To reduce the possibility of disease resistance to Rovral[®], the alternation with other fungicides is recommended.

PLUM/PRUNE - CONTROL OF BROWN ROT

RATE: 1.5 kg/ha of Rovral[®].

TIMING: To protect the fruit use Rovral[®] in a cover spray program with application at 7-14 day intervals throughout the season. Use the short interval under wet conditions close to harvest. Observe a 1 day PHI. Apply by ground. To reduce the possibility of disease resistance to Rovral[®], alternation with other fungicides is recommended.

RASPBERRY - CONTROL OF GREY MOLD FRUIT ROT

RATE: 2 kg/ha of Rovral[®] in 1000 litres of water per hectare.

TIMING: Apply a maximum of 8 applications per year. Begin applications when flower buds are visible in the crown and repeat at 7-10 day intervals. Reduce intervals during rainy periods. Apply by ground. Observe a 1 day PHI. To reduce the possibility of disease resistance to Rovral[®], alternation with other fungicides is recommended.

STRAWBERRY - CONTROL OF BOTRYTIS FRUIT ROT AND SUPPRESSION OF PENICILLIUM

RATE: 2 kg/ha of Rovral[®] in at least 1000 litres of water per hectare. Good coverage of the plants is essential. The addition of a non-ionic adjuvant registered for this purpose will improve performance.

TIMING: Begin application when flower buds are visible in the crown and repeat at 7-10 day intervals. Reduce interval during rainy periods. To reduce the possibility of disease resistance to Rovral[®], the alternation with other fungicides is recommended. Leave 1 day from last spray to harvest.

Booklet Label

CABBAGE (STORED) - CONTROL OF ALTERNARIA

RATE: 3 kg/ha of Rovral®.

TIMING: Make 1 application 7-14 days before harvest. Apply by ground; boom or airblast sprayer. Observe a PHI of 7 days.

CAULIFLOWER - CONTROL OF ALTERNARIA

RATE: 3 kg/ha of Rovral®.

TIMING: Make 1 application 1-2 days prior to tying. Apply by ground; boom or airblast sprayer. Observe a PHI of 5 days.

CUCUMBER (GREENHOUSE) - CONTROL OF GUMMEY STEM BLIGHT

RATE: 100g Rovral®/100 L water applied at 1000-2000 L water per hectare. Spray to run-off ensuring good coverage of the plants, particularly stem nodes, immediately following any pruning.

TIMING: Begin application at the first sign of disease. Spray at weekly intervals or as required for control. Observe a PHI of 2 days.

CUCUMBER (GREENHOUSE) - CONTROL OF BOTRYTIS GREY MOLD

RATE: 100g Rovral®/100 L water per hectare. Spray to run-off ensuring good **coverage of the plants particularly stem nodes immediately following any pruning.**

TIMING: Begin application at the first sign of disease. Spray at weekly intervals or as required for control. At least 2 days must elapse between treatment and picking. To reduce the possibility of disease resistance, Rovral® must be alternated with other fungicides.

GARLIC - CONTROL OF GREEN MOLD (*Penicillium corymbiferum*)

RATE: 4g/L Rovral® as a seed dip.

TIMING: Dip garlic seeds for 30 minutes in a suspension of Rovral® and water. Make one application in the fall, immediately before planting.

GINSENG - CONTROL OF ALTERNARIA LEAF BLIGHT

RATE: 1.1 Kg/ha Rovral® in at least 2000 L of water per hectare.

TIMING: Apply when the disease first appears and follow with two subsequent sprays at monthly intervals as part of a spray programme. Do not apply more than three applications per season; the Rovral® applications should be alternated with other suitable fungicides. Do not apply closer than 30 days to harvest.

Booklet Label

LETTUCE (FIELD)- CONTROL OF GREY MOLD

RATE: 1.5 kg/ha of Rovral® in 300-500 L water per hectare.

TIMING: Apply a maximum of 4 applications per year at 7 day intervals between sprays. Apply by ground; boom sprayer. Observe a PHI of 4 days.

LETTUCE (GREENHOUSE) - CONTROL OF GREY MOLD/SCLEROTINIA DROP

RATE: 2 kg/ha of Rovral® in 2000 L water per hectare. Use a high volume sprayer.

TIMING: Apply a maximum of 4 applications per year with the first application at the 3-leaf stage and then every 7-10 days if conditions continue to favour disease development. Observe a PHI of 14 days.

ONION (DRY BULB) - CONTROL OF BOTRYTIS LEAF BLIGHT

RATE: 1.5 kg/ha Rovral® applied in 300-500 litres of water. The addition of a non-ionic adjuvant will improve performance.

TIMING: Begin applications about mid-June or at the first sign of leaf spotting and repeat at 7-10 day intervals.

Do not apply Rovral® within 15 days of harvest. To reduce the possibility of disease resistance to Rovral®, the alternation with other fungicides is recommended.

ONIONS (DRY BULB) - CONTROL OF BOTRYTIS LEAF BLIGHT PLUS DOWNY MILDEW

Manzate DF Fungicide

RATE: Use 0.75 kg/ha Rovral® tank-mixed with the recommended rate of Maneb 80W or Manzate 200 for downy mildew control (consult the Maneb 80W or Manzate 200 label for dose rates, precautions and limitations). Apply in 300-500 litres of water per hectare. The addition of a non-ionic adjuvant will improve performance.

Manzate DF Fungicide or Manzate Pro-Stick Fungicide

TIMING: Apply this tank-mix as a preventative treatment. Application can begin mid-June or when conditions are favourable for disease infection. Repeat applications at 7-10 day intervals. DO NOT apply Rovral® within 15 days of harvest.

TOMATO (GREENHOUSE) - CONTROL OF BOTRYTIS GREY MOLD

RATE: 100 g Rovral®/100 L water. Spray to run-off ensuring good coverage of the plants, particularly stem nodes immediately following any pruning.

TIMING: Treat when the second truss is in flower or at the onset of the disease. Repeat treatment at weekly intervals or when conditions are favourable.

At least 2 days must elapse between treatment and picking. To reduce the possibility of disease resistance, Rovral® must be alternated with other fungicides.

Booklet Label

CANOLA - CONTROL OF SCLEROTINIA STEM ROT

RATE: 1-1.5 kg/ha Rovral[®] as an aerial application in at least 45 litres of water per hectare. The use of the higher rate is recommended for fields with a history of heavy disease pressure or for dense crop stands. Good coverage of the plants is essential.

The addition of a non-ionic adjuvant will improve performance.

TIMING: Apply Rovral[®] when 20-30% of the blooms have opened.

DRY COMMON BEANS (WHITE & KIDNEY) - CONTROL OF SCLEROTINIA WHITE MOLD

RATE: 1-1.5 kg/ha Rovral[®]. The use of the higher rate is recommended for fields with a history of heavy disease pressure or for dense crop stands. For ground application use 300 litres of water per hectare. For aerial application, use at least 45 litres of water per hectare. Good coverage of the foliage and stems is essential. To ensure thorough coverage for ground applications, a combination of overhead and drop nozzles should be used at a minimum pressure of 400 kPa. The addition of a non-ionic adjuvant will improve performance.

TIMING: Apply Rovral[®] as a preventative treatment when 25-75% of the blooms have opened. Treatment prior to the presence of disease is preferable, but Rovral[®] is still effective if application is made at the initial sign of disease infection, when less than 5% of the plants are showing white mold.

SNAP BEANS - CONTROL OF SCLEROTINIA WHITE MOLD AND BOTRYTIS POD ROT

RATE: 1.5 kg/ha of Rovral[®] in 300 litres of water per hectare for ground applications and at least 45 litres of water for aerial application. Good coverage of the plants is essential. The addition of a non-ionic adjuvant registered for this purpose will improve performance.

TIMING: Control of Sclerotinia white mold - apply Rovral[®] when 50% of the blooms have opened. Control of Botrytis pod rot (normally only a problem in British Columbia) - apply Rovral[®] twice, when 30% and again when 50% of the blooms have opened.

CARROT SEED - CONTROL OF SEED BORNE ALTERNARIA

RATE: 521 mL of Rovral[®]/100 kg seed

DIRECTIONS: For the importation of treated seed into Canada; NO SEED IS TO BE TREATED IN CANADA:

Wear protective gloves and overalls when handling treated seeds. When augering or handing treated seed work in a well ventilated area and wear a suitable respirator.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS (BELOW): The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Bayer CropScience and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Bayer CropScience itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below. Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Bayer CropScience harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

LEEK – CONTROL OF *BOTRYTIS* LEAF BLIGHT

Pest: For control of Botrytis leaf blight

Rate: 1.5 kg/ha of Rovral® applied in 300-500 litres of water as a foliar spray.

Timing: Apply a maximum of 4 applications per year with the first application at the appearance of the disease and /or when conditions are favourable to the disease prevail. Repeat applications should be made at 7-10 day intervals. Observe a PHI of 15 days.

TURF AND ORNAMENTALS

TURF

Rovral® is effective either as a preventative or curative spray. Ensure the area to be treated is covered uniformly. Apply the required amount of Rovral® in not less than 10 litres per 100 m². When treating golf greens, always treat aprons and fairway approaches. Avoid mowing for ten hours after application.

DOLLAR SPOT (*Sclerotinia homeocarpa*)

Apply Rovral® at the first sign of disease and repeat at recommended intervals as long as required. Use either 30g per 100 m² at 14-day intervals or 45 g per 100 m² at 21 day intervals.

BROWN PATCH (*Rhizoctonia solani*)

Apply Rovral® at 60 to 125 g per 100 m², beginning when the disease is first observed and continue at 14 to 21 day intervals throughout the season or as needed. Under severe disease pressure use the higher rate and the shorter interval.

LEAF SPOT, MELTING OUT (*Helminthosporium* spp.)

Apply Rovral® at 60 g per 100 m² when the disease first appears. Repeat 2 or 3 times in the early spring at 14 -21 day intervals or as needed.

Booklet Label

GREY AND PINK SNOW MOLD (*Typhula* spp. and *Fusarium nivale*)

To suppress the active stages of the disease during spring or fall, apply Rovral® at 180g per 100 m². For winter protection, Rovral Green® is recommended.

ORNAMENTALS - CONTROL OF BOTRYTIS spp.

RATE: 10g Rovral® per 10 litres of water. Apply to the point of run-off, ensuring good coverage of the plants.

TIMING: Treat at the onset of Botrytis and repeat at 21 day intervals, or as necessary.

To reduce the possibility of disease resistance, Rovral® **must** be alternated with other fungicides.

The following species have been safely treated:

Adiantum spp	Hypoestes sanguinolenta
Apehlandra squarrosa	Impatiens spp
Aralia elegantissima	Iresine herbstii
Aralia sieboldii	Kalanchoe spp
Asparagus sprengeri	Maranta spp
Asparagus plumosus	Monstera deliciosa borsigiana
Azalea spp	Neanthe bella
Begonia rex var. Fireglow	Nepeta (ginger plant)
Chrysanthemum spp	Nephrolepis spp
(all year round)	Pelargonium spp
Chlorophytum spp	Peperomia caperata
Cineraria spp	Peperomia hederifolia
Ciccus antarctica	Peperomia magnoliifolia
Croton spp var. Bravo	Philodendron scandens
Cyclamen spp	Pilea cadierei
Dracaena spp var. Rededge	Poinsettia spp
Episcia cupreata	Primula spp
Euonymus spp	Rhoicissus spp
Euphorbia splendens	Ruellia makoyana
Ficus spp	Saintpaulia ionantha**
Fittonia spp	Saxifraga stolonifera
Fuchsia spp	Senecio macroglossus variegatum
Geranium (Zonal)	Setcreasea purpurea
Gesneria spp	Sinningia spp (Gloxinia)
Gynura sarmentosa	Solanum capsicastrum
Hedera spp	Rosa samantha

** Spraying Saintpaulia flowers may result in petal scorch. Tepid water should be used and the wet plants protected from direct sunlight.

Tolerance of other species should be checked on a few plants before making any large scale applications.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.