



Overview

Fungicides

Herbicides

Alion

Betamix B

Buctril M

Centurion

Converge

Excel Super

Fusion

Ignite

Infinity

Liberty 150

Liberty 200

Nortron SC

Option Liquid

Pardner

Home » Products » Herbicides » Puma Advance

## Puma Advance

### Overview

### Labels & MSDS

### Application Guidelines

#### Overview

- Affordable wild oat control offers excellent return on investment
- Proven control of wild oats, barnyard grass, yellow foxtail and green foxtail
- **Crop safe** – Improved crop safety over Puma® Super especially in barley
- **Flexible** – Mixes well with Infinity® and over 25 other broadleaf herbicides
- **Wide window of application** – 1 to 6 leaves on main stem plus 3 tillers
- Registered for use in both wheat (spring and durum) and barley
- Move safely from wheat to barley without having to clean out your sprayer

### Search

Region: Ontario

### MyBayer

>> Forgot Password

[Puma Advance](#)  
[Select](#)  
[Sencor](#)  
[Thumper](#)  
[Tundra](#)  
[Varro](#)  
[Velocity m3](#)  
[Vios G3](#)  
[BigBoys](#)  
[Insecticides](#)  
[Other](#)  
[Seed](#)  
[Seed Treatments](#)

#### Bayer Links

[Bayer Global](#)  
[Bayer CropScience Global](#)

- No re-cropping restrictions
- The only advanced fenoxaprop formulation to offer increased crop safety and improved wild oat performance
- Registered for both ground and aerial application

#### Application Tips

Rates	
Barley and wheat	For control of wild oats, green foxtail, yellow foxtail and barnyard grass in barley and wheat, mix Puma Advance at 0.412 L/ac.

Timing
For wheat and barley, the crop stage is between 1 to 6 leaves on main stem plus 3 tillers. Apply at the 3 to 4 leaf stage to achieve maximum crop tolerance and weed control.

Packaging
<ul style="list-style-type: none"> <li>• 8.25 L jug = 20 acres (one 2 x 8.5 L case = 40 acres)</li> <li>• 123.75 L BigBoys shuttle = 300 acres</li> <li>• 412.5 L BigBoys tote = 1,000 acres</li> </ul>

Water Volume
<ul style="list-style-type: none"> <li>• Ground – minimum of 6 US gal./ac. (22 L/ac.)</li> <li>• Aerial – minimum of 4 US gal./ac. (14 L/ac.)</li> </ul>

Rainfastness
One hour after application

Residue and Grazing
<p>Do not:</p> <ul style="list-style-type: none"> <li>• Do not graze the treated crops or cut for hay within 25 days of Puma Advance herbicide application, or harvest for grain within 65 days of application</li> </ul>

#### Find My Local Bayer Rep



Don't know your local Bayer CropScience representative?



- Do not graze the treated perennial ryegrass or cut for straw within 65 days of application
- Do not graze the treated meadow brome grass or cut for hay within 25 days of application

---

Last updated:  
May 22, 2015

- » [General Terms of Use](#)
- » [Privacy Statement](#)
- » [About](#)
- » [General Terms of Use](#)

---

[Social Media](#)  
[Print](#)  
[Bookmark](#)