

GROUP	3	FUNGICIDE
-------	---	-----------

FUNGINEX® DC Fungicide

Emulsifiable Concentrate

AGRICULTURAL

GUARANTEE: Triforine190 g/L

REGISTRATION NO. 27686

PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

DANGER - EYE IRRITANT and SKIN IRRITANT

IN CASE OF EMERGENCY INVOLVING A POISONING OR MAJOR SPILL
CALL

1-866-336-2983 OR 1-703-527-3887

NET CONTENTS: 0.5 L to 10 L

REGISTRANT
SUMMIT AGRO NORTH AMERICA HOLDING CORPORATION
600 THIRD AVENUE
NEW YORK NY 10016-2001
UNITED STATES

Canadian Representative
Engage Agro Corporation
1030 Gordon St.
Guelph, ON
N1G 4X5
519-826-7878

FUNGINEX® Trademark registered by Sumitomo Corporation, Japan, under license to Summit Agro (USA) Corp.

DIRECTIONS FOR USE

BLUEBERRY (LOWBUSH & HIGHBUSH)

- a) BRITISH COLUMBIA - for control of mummyberry, apply 3 litres in 1,000 L of water/ha.
- b) EASTERN CANADA - for control of mummyberry, apply 1.7 to 3 litres per hectare.

Apply in the spring at bud break and repeat 10 - 14 days later. Apply a third spray at early bloom and repeat 10 - 14 days later. Thoroughly wet all buds and shoots. Do not apply later than 60 days before harvest. Consult local extension service for proper application timing.

NOTE: (EASTERN CANADA ONLY): Do not make more than three applications from leaf-bud break to pink-bud stage. Application of FUNGINEX DC Fungicide during or beyond early bloom may result in fruit russetting.

Ground application only, apply in 1,000 L of water/ha.

DO NOT APPLY BY AIR

CRANBERRY (B.C. ONLY): For control of cottonball apply 3 L per hectare in 1,000 - 1,500 L of water in the spring at leaf bud-break. Repeat 10 - 14 days later. Apply third spray at early bloom and repeat 10-14 days later. Thoroughly wet all buds and shoots. Do not apply later than 60 days before harvest. Consult local extension service for proper application timing. **Do not apply by air.**

SASKATOON BERRIES: For the control of Saskatoon-juniper rust (*Gymnosporangium* spp.) and Entomosporium leaf and berry spot, apply 3 L/ha in a water volume (e.g., 1000 L) sufficient to ensure thorough wetting - to the point where the solution drips from the plants. Actual orchard area covered will vary, depending on size of bushes and density of planting. **Do not apply by air.**

TIMING: A single application should be made between flower bud break and the white tip stage. Do not apply later than 60 days before harvest.

PEACHES, CHERRIES, PLUMS, PRUNES: For control of the blossom blight stage of brown rot, apply 750 mL of FUNGINEX DC Fungicide per 1,000 L of water or 2.5 L per hectare. Make 2 - 3 applications from early to full bloom stage, depending on the length of the bloom period and conditions favouring brown rot. Follow provincial recommendations for proper timings. **Do not apply by air.**

APPLE NURSERY STOCK AND NONBEARING APPLE TREES: For control of powdery mildew, apply 2.5 L FUNGINEX DC Fungicide per hectare by airblast sprayer at tight cluster, pink and petal fall stages. May be applied 2 weeks after petal fall (mid-

summer) if necessary. Make a maximum of 3-5 applications per season. **Do not apply by air.**

ROSES AND ORNAMENTALS: For control of powdery mildew and black spot, mix 1 L per 1,000 L of water. Spray to thoroughly cover all plant surfaces. **Do not apply by air.**

DO NOT use in greenhouses.

The maximum number of applications per year allowed on roses and ornamentals is 3.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

NOTE: Do not let spray mixture stand in the tank overnight.

ENVIRONMENTAL HAZARDS:

TOXIC to aquatic organisms, birds and small wild mammals. Observe buffer zones specified under **DIRECTIONS FOR USE.**

To minimize surface water contamination when used on cranberries, all effluent water must be impounded and released only when levels of the active ingredient are <1100 µg a.i./L.

The use of the chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g., sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Buffer zones:

Use of the following spray methods or equipment DO NOT require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop	Buffer Zones (metres) Required for the Protection of Aquatic Habitat at Water Depths of:	
		Less than 1 m	Greater than 1 m
Field sprayer	Lowbush blueberries, cranberries, apple nursery stocks and nonbearing apple trees	1	0
Airblast	Cranberries, highbush blueberries, apple nursery stocks and nonbearing apple trees	1	0

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that FUNGINEX DC Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to FUNGINEX DC Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of FUNGINEX DC Fungicide or other Group 3 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group when such use is permitted.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.

- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact Engage Agro Corp at 1-866-613-3336.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed.

Severe irritant to eyes. DO NOT get in eyes.

Causes skin irritation and DO NOT get on skin.

Avoid inhalation of spray mist.

Wash thoroughly with soap and water after using and before eating or smoking.

Wear chemical-resistant coveralls over long-sleeve shirt and long pants, chemical-resistant gloves and goggles during mixing, loading, application, clean up and repair.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Avoid contamination of food and feed and domestic or irrigation water supplies, lakes, streams and ponds.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at: www.croplife.ca.

DO NOT APPLY BY AIR

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE

1. Store above 5° Celsius.
2. Store in original, tightly closed container.
3. Do not ship or store near food, feed, seed and fertilizers.
4. Store in cool, dry, locked, well-ventilated area without floor drain.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the **PEST CONTROL PRODUCTS ACT** to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.