

GROUP

4

INSECTICIDE

QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE

FOR SYSTEMIC CONTROL OF THE LARVAE OF EUROPEAN CHAFER, JAPANESE
BEETLE AND BLACK TURFGRASS ATAENIUS BEETLE AND SUPPRESSION OF
EUROPEAN CRANE FLY LARVAE TURFGRASS INCLUDING SOD FARMS

COMMERCIAL

GUARANTEE: Imidacloprid 0.5%

REGISTRATION NO. 29185 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

CAUTION



POISON

EYE IRRITANT

NET CONTENTS: 5-50 kg

Makhteshim Agan of North America Inc.
4515 Falls of Neuse Road, Suite 300
Raleigh, NC 27609
919 256-9300

DIRECTIONS FOR USE

Read the entire DIRECTIONS FOR USE AND GENERAL INFORMATION before using QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE.

GENERAL INFORMATION

APPLICATION TO TURFGRASS

QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE can be used for the control of soil inhabiting life stages of European chafer beetle, Japanese beetle, and Black Turfgrass Ataenius beetle and the suppression of European crane fly larvae (leather jackets). QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE can be used as directed on turfgrass in sites such as home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms.

The active ingredient in QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE has sufficient residual activity so that applications can be made preceding the egg-laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg-laying period. The need for an application can be based on monitoring of the site, previous records or experiences, current season adult trapping or other methods. Applications should be made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall (5-10 mm) to move the active ingredient through the thatch. Avoid overwatering (more than 20 mm). Avoid runoff or puddling of irrigation water following application.

The treated turf area must be in such a condition that the rainfall or irrigation water will penetrate vertically into the soil profile, which will ensure an adequate distribution of the active ingredient. Avoid application of QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE to turfgrass areas which are waterlogged or saturated, which will not allow penetration of the active ingredient into the plant root zone.

Apply QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE ONLY ONCE per year as directed by this label.

APPLICATION EQUIPMENT

Apply QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE uniformly over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately. **DO NOT APPLY BY AIR.**

SPREADER SETTINGS FOR QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE. The chart below can be used as guide for establishing an accurate rate of QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE for several commonly used application equipment. Application rates will vary depending upon speed, equipment wear and condition of spreader. Carefully calibrate application equipment prior to commercial use of this product.

SPREADER	SPEED (km/h)	SETTINGS
		56 kg/ha
Scotts Acupro 2000	5	G ½
Scotts SR 2000	5	G ½
Scotts R 8	5	H
Lesco 6000	5	B ½
Lesco Commercial Plus (Model 705699)	5	7

RECOMMENDED APPLICATIONS			
CROP	PEST	DOSAGE QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE	REMARKS
Lawns: Home, business and office complexes, shopping complexes, multi-family residential complexes, airports, cemeteries, parks, playgrounds, athletic fields, golf courses, sod farms.	For Control of the Larval Stage: Japanese beetle, European chafer beetle, Black turfgrass ateanius	56 kg/ha or 560 g per 100 m ²	For optimum control, make application prior to egg hatch of the target pest. * For suppression of early fall larval stages of crane fly (larvae known as leatherjackets) apply QUALI-PRO IMIDACLOPRID 0.5 Granular Insecticide to turfgrass areas known to be infected with leatherjackets. Application timing should commence when adult crane flies take flight in August and September and begin laying eggs.
	For Suppression of the Larval Stage: European Crane Fly Larvae*		
<p>Be sure to read “APPLICATION EQUIPMENT” Section of this label. Consult your local Agricultural (Horticultural) Advisor for more specific information regarding timing of application. In Halifax, Nova Scotia, consult Canadian Food Inspection Agency personnel for more specific information regarding monitoring and timing of application. NOTE: For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. On golf courses, irrigate treated areas following application. Avoid mowing turf or lawn area until after irrigation or rainfall has occurred so that uniformity of application will not be affected. Apply QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE only once per year as directed by this label. If QUALI-PRO IMIDACLOPRID 0.5 GRANULAR Insecticide was applied to a turfgrass area earlier in the summer season for Japanese beetle or European chafer or Black turfgrass ateanius, do not re-apply for leatherjacket suppression in the same year</p>			

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE contains a Group 4 insecticide. Any insect population may contain individuals naturally resistant to QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE and other Group 4 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE or other Group 4 insecticides with different groups that control the same pests in a field.
- Insecticide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Before applying QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE, correctly identify the pest and ensure economic and agronomic thresholds are met as recommended by local provincial or IPM specialists.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Makhteshim Agan of North America Inc. via internet at www.manainc.com or phone 1-919 256-9300.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled. May cause irritation of the respiratory tract. Avoid inhaling dust. Keep unprotected persons, children and pets out of the treated areas until irrigation has dried. Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt, long pants, and chemical-resistant gloves and a suitable dust-filtering respirator approved by NIOSH/MSHA with any R, P, or HE filter during handling, loading, applying, clean-up and repair. Wash thoroughly with soap and water after handling.

RESTRICTIONS

Do not graze treated areas or use clippings from treated areas for feed or forage. Do not plant any food crop within one year of a treatment with QUALI-PRO IMIDACLOPRID 0.5 GRANULAR INSECTICIDE.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, ponds or other aquatic systems.

Do not apply to terrains where there is a potential for surface run-off to enter aquatic systems.

Do not mix, load or clean equipment within 30 metres of wellheads or aquatic systems. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to aquatic invertebrates. Apply this product only in accordance with this label.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal.

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.