

[Bag Label]

GROUP

2

HERBICIDE

ULTIM® 75 DF HERBICIDE

DRY FLOWABLE

FOR SALE FOR USE ON FIELD CORN IN EASTERN CANADA AND MANITOBA, AND LOWBUSH
BLUEBERRIES IN EASTERN CANADA ONLY

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE: Rimsulfuron 37.5%
Nicosulfuron 37.5%

REGISTRATION NO. 24736 PEST CONTROL PRODUCTS ACT

CAUTION - SKIN IRRITANT
WARNING - EYE IRRITANT

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

NET CONTENTS: 134.8 GRAMS
(4 x 33.7 gram water soluble bags)

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Potential skin sensitizer.
- Avoid breathing spray mist.
- Avoid contact with skin, eyes and clothing.
- Do not contaminate any body of water.
- DO NOT APPLY BY AIR

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact 1-866-375-4648 or www.croplife.ca/.

FIRST AID:**EYE CONTACT:**

Flush eyes immediately with plenty of clean water for 15 minutes. Contact a physician, or poison control centre IMMEDIATELY.

SKIN CONTACT:

Wash skin with plenty of soap and water. Remove and wash contaminated clothing before reuse.

INGESTION:

No specific intervention is indicated as compound is not likely to be hazardous by ingestion. Consult physician or a poison control centre IMMEDIATELY

INHALATION:

No specific intervention is indicated as the compound is not likely to be hazardous by inhalation. Consult a physician or poison control centre IMMEDIATELY.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Keep water soluble bags in original outer bag or outer container away from moisture.

DISPOSAL:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For mylar bag containers:

1. Make the empty bag unsuitable for further use.
2. Dispose of the bag in accordance with provincial requirements.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe on any Canadian patent.

[DFU Pamphlet]

GROUP	2	HERBICIDE
-------	---	-----------

ULTIM® 75 DF HERBICIDE

DRY FLOWABLE

FOR SALE FOR USE ON FIELD CORN IN EASTERN CANADA AND MANITOBA, AND LOWBUSH
BLUEBERRIES IN EASTERN CANADA ONLY

AGRICULTURAL

READ THE LABEL AND THIS BOOKLET BEFORE USING

GUARANTEE: Rimsulfuron 37.5%
Nicosulfuron 37.5%

REGISTRATION NO. 24736 PEST CONTROL PRODUCTS ACT

CAUTION - SKIN IRRITANT
WARNING - EYE IRRITANT

E. I. DU PONT CANADA COMPANY
AGRICULTURAL PRODUCTS
P.O. BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Potential skin sensitizer.
- Avoid breathing spray mist.
- Avoid contact with skin, eyes and clothing.
- Do not contaminate any body of water.
- **DO NOT APPLY BY AIR.**
- Do not sell water soluble bags separately

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact 1-866-375-4648 or www.croplife.ca/.

FIRST AID:

EYE CONTACT:

Flush eyes immediately with plenty of clean water for 15 minutes. Contact a physician, or poison control centre IMMEDIATELY.

SKIN CONTACT:

Wash skin with plenty of soap and water. Remove and wash contaminated clothing before reuse.

INGESTION:

No specific intervention is indicated as compound is not likely to be hazardous by ingestion. Consult physician or a poison control centre IMMEDIATELY.

INHALATION:

No specific intervention is indicated as the compound is not likely to be hazardous by inhalation. Consult a physician or poison control centre IMMEDIATELY.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-800-441-3637 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

APPLICATION PRECAUTIONS:

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following: Do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water. Keep from contact with fertilizers, insecticides, fungicides and seeds.
- APPLY ULTIM® 75 DF HERBICIDE WITH A RECOMMENDED SURFACTANT.
- DO NOT APPLY BY AIR.
- Do not handle pesticides with bare hands. Chemical-resistant gloves significantly reduce hand exposure. Wear gloves for mixing/loading/clean-up operations and when making sprayer and nozzle repairs and adjustments. Do not use leather or cloth gloves.
- Carefully observe sprayer cleanup instructions, as spray tank residue may damage crops other than corn.

GENERAL INFORMATION:

ULTIM® 75 DF Herbicide is a dry flowable granular formulation to be mixed in water and applied postemergence to field corn in Eastern Canada for control of quackgrass, annual grasses, and redroot pigweed. ULTIM® 75 DF Herbicide is pre-packaged in water soluble bags; each 33.7 gram water-soluble bag treats 1 hectare. Use gloves when handling the water-soluble bags. Reseal unused water-soluble bags into outer bag or outer container when not in use. Keep water-soluble bags away from excess moisture or physical damage during storage and prior to mixing in the spray tank. Do not reuse outer bag or outer container.

ULTIM® 75 DF Herbicide is non-corrosive, non-flammable, non-volatile, and does not freeze. ULTIM® 75 DF Herbicide rapidly stops growth of susceptible species; typical symptoms usually appear within 5-7 days, but may not be noticeable for 2-3 weeks after application, depending on growing conditions and weed susceptibility. Warm, moist conditions following application promote the activity of ULTIM® 75 DF Herbicide, while cool and/or dry conditions may reduce or delay activity. Weeds hardened off by cold weather or drought stress may not be controlled.

Weed control may be reduced if the corn canopy has closed in over the weeds, intercepting the spray.

ULTIM® 75 DF Herbicide is a contact herbicide, and will not give residual control of grass or broadleaf weed seedlings that may germinate after application.

DIRECTIONS FOR USE:

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

- A rapid fluctuation in temperature (greater than 20 °C difference within 24-36 hours) will stress the corn crop. For maximum crop safety, allow 48-72 hours for the corn to acclimatize before spraying ULTIM® 75 DF Herbicide.
- For ULTIM® 75 DF Herbicide used alone or in a tankmix, apply ONLY when the temperature in the 24 hours before AND after application ranges between 5°C and 28 °C. Temperatures beyond this range increase the potential for crop injury. Separate applications of ULTIM® 75 DF Herbicide followed by a broadleaf herbicide (minimum 12 hours later) will reduce the potential for injury.
- Ensure that the boom is set at the proper height in relation to the corn plants to apply ULTIM® 75 DF Herbicide accurately and uniformly, and to AVOID excessive application into the corn whorl.
- WARNING: Crop injury may result if application is made to corn that has been stressed by abnormally hot, humid or cold weather conditions, frost, low fertility, drought, water saturated soil, compacted soil, previous pesticide applications, disease or insect damage. If corn has been injured by frost, wait 48-72 hours before applying ULTIM® 75 DF Herbicide.
- When tank mixing ULTIM® 75 DF Herbicide with a broadleaf herbicide, always read the broadleaf herbicide label before using and follow all precautions.
- ULTIM® 75 DF Herbicide rapidly stops growth of susceptible species; typical symptoms usually appear within 5-7 days, but may not be noticeable for 2-3 weeks after application, depending on growing conditions and weed susceptibility.
- Warm, moist conditions following application promote the activity of ULTIM® 75 DF Herbicide, while cool and/or dry conditions may reduce or delay activity. Weeds hardened off by cold weather or drought stress may not be controlled.
- ULTIM® 75 DF Herbicide is a contact herbicide, and will not give residual control of grass or broadleaf weed seedlings that may germinate after application.
- Weed control may be reduced if the corn canopy has closed in over the weeds, intercepting the spray.

CORN HYBRID SENSITIVITY:

Corn hybrids with corn-heat-unit (CHU) ratings of 2500 or less have shown some sensitivity to ULTIM® 75 DF Herbicide. DO NOT USE ULTIM® 75 DF HERBICIDE ON CORN HYBRIDS WITH CHU RATINGS OF 2500 OR LESS OR IN GEOGRAPHIC REGIONS HAVING 2500 OR LESS AVERAGE SEASONAL CORN HEAT UNITS.

Corn hybrids of 2550 CHU or greater that have shown considerable crop injury from applications of ULTIM® 75 DF Herbicide are listed below. ULTIM® 75 DF Herbicide should not be sprayed for weed control in fields planted with these hybrids.

PIONEER® BRAND 3897	MAX 78
PIONEER® BRAND 3936	G-4034
PIONEER® BRAND 3941	G-4120
PICKSEED® 2525	G-4160

Do not use on sweet corn. Do not use on seed corn or popcorn.

ULTIM® 75 DF HERBICIDE USED ALONE

SPRAY ADJUVANTS:


ULTIM® 75 DF Herbicide must be applied with a recommended non-ionic surfactant, either CITOWETT* PLUS, AGRAL* 90 or AG-SURF* at 2 L per 1000 L spray solution (0.2% v/v) or SIDEKICK* at 2.5 L per 1000 L spray solution (0.25% v/v).

RATES AND TIMING:

Apply one water soluble bag of ULTIM® 75 DF Herbicide per hectare as a broadcast spray, with a recommended surfactant to corn and weeds according to the following chart:

Crop or Weed	Leaf Staging at Application
Corn	1-6 leaves (4 visible collars or 30 cm in height - leaf extended) See diagram below
Annual Grasses	1-6 leaves (up to early tillering, 2 - two leaf tillers)
Long-spined Sandbur	3-5 leaves
Quackgrass	3-6 leaves (10-20 cm in height - leaf extended)
Redroot Pigweed	2-6 leaves (10 cm tall or across)

CORN LEAF STAGING



Apply ULTIM® 75 DF Herbicide from the 1 to 6-leaf stage (4 visible collars or 30 cm in height - leaf extended).

The coleoptile (short, blunt leaf) is counted as the first leaf. A corn leaf is counted as a full leaf when the next leaf is visible in the corn whorl.

If using a tank mixture with another herbicide, refer to the section on tank mixtures below for correct timing.

Applications of ULTIM® 75 DF Herbicide after the 6-leaf stage of corn (30 cm in height leaf extended) must be delayed until after the 8-leaf stage and before silking, when a height differential can be established between the corn and target weeds. When drop piping, ensure that the crop stand is UNIFORM so that all the spray solution is directed below the canopy to keep the spray out of the corn whorl. Application at the later stage must be directed below the corn canopy to keep the spray out of the corn whorl.

WEEDS CONTROLLED BY ONE WATER SOLUBLE BAG/HA:

Barnyard Grass	Proso Millet ²	Yellow Foxtail [†]
Fall Panicum	Quackgrass ¹	Old Witch Grass
Green Foxtail	Redroot Pigweed (including triazine resistant)	
Long-spined sandbur		

[†] Suppression only

NOTE: Weeds that emerge after application of ULTIM® 75 DF Herbicide will not be controlled. Application should be made when the majority of weeds have emerged.

Annual grasses and redroot pigweed are most sensitive when small and actively growing.

Early crop establishment and a good crop stand (minimum of 50,000 plants per hectare) are important in providing competition for weeds and for effective postemergence control of quackgrass and annual grasses.

Specific Weed Control Recommendations:

1. Quackgrass - Apply ULTIM® 75 DF Herbicide with a recommended surfactant postemergence when the majority of the quackgrass shoots are actively growing and in the 3 to 6-leaf stage (10 to 20 cm in height - leaf extended). If the correct leaf stage of quackgrass does not coincide with the correct leaf stage of broadleaf weeds (see tank mix section of this label), separate applications of ULTIM® 75 DF Herbicide and a broadleaf herbicide are required. Annual grasses not emerged at the time of ULTIM® 75 DF Herbicide application to quackgrass will not be controlled.
2. Proso Millet - Apply ULTIM® 75 DF Herbicide with a recommended surfactant postemergence when the majority of proso millet has emerged and before the largest proso millet plants reach the multi-tillered (3 to 4 tillers) stage. If, at correct growth stage for proso millet, corn is beyond the 6- leaf stage, application of ULTIM® 75 DF Herbicide must be delayed until after the 8-leaf stage of corn and before silking, when a height differential can be established between the corn and proso millet. When drop piping, ensure that the crop stand is uniform so that all the spray solution is directed below the canopy to keep the spray out of the corn whorl. Reduced control of proso millet may result from this timing of application due to the advanced stage of the weed plants.

ULTIM® 75 DF HERBICIDE TANK MIXES

ULTIM® 75 DF HERBICIDE + BANVEL* HERBICIDE

ULTIM® 75 DF Herbicide may be tank mixed with BANVEL* Herbicide to control annual grasses, quackgrass and many broadleaf weeds. Tank mix one water-soluble bag/ha of ULTIM® 75 DF Herbicide and 600 mL/ha of BANVEL* Herbicide with a non-ionic surfactant at 0.2% v/v (2 L/1000 L) and apply postemergence to corn up to the 6-leaf stage (4 visible collars or 30 cm in height - leaf extended).

This tank mixture will control the weeds listed for control by ULTIM® 75 DF Herbicide alone, plus the weeds listed on the BANVEL* Herbicide label. Refer to the "ULTIM® 75 DF HERBICIDE USED ALONE" section of this label for more application instructions and restrictions on the use of ULTIM® 75 DF Herbicide. Refer to the BANVEL* Herbicide label for weeds controlled by BANVEL* Herbicide and consult the label for additional application instructions and use precautions.

The tank mix of ULTIM® 75 DF Herbicide and BANVEL* Herbicide must not be applied after the 6-leaf stage (4 visible collars or 30 cm in height - leaf extended) of corn.

ULTIM® 75 DF HERBICIDE + BANVEL* II HERBICIDE

ULTIM® may be tank mixed with BANVEL* II Herbicide to control annual grasses, quackgrass and many broadleaf weeds. Tank mix one water-soluble bag/ha of ULTIM® 75 DF Herbicide and 600 mL/ha of BANVEL* II Herbicide with a non-ionic surfactant at 0.2% v/v (2 L/1000 L) and apply postemergence to corn from the spike to the 6-leaf stage (4 visible collars or 30 cm in height - leaf extended).

This tank mixture will control the weeds listed for control by ULTIM® 75 DF Herbicide alone, plus the weeds listed on the BANVEL* II Herbicide label. Refer to the "ULTIM® 75 DF HERBICIDE USED ALONE" section of this label for more application instructions and restrictions on the use of ULTIM® 75 DF Herbicide. Refer to the BANVEL* II Herbicide label for weeds controlled by BANVEL* II Herbicide and for additional application instructions and use precautions.

The tank mix of ULTIM® 75 DF Herbicide and BANVEL* II Herbicide should be applied at weed stage of emergence to 6-leaf. Do not apply after 6-leaf (4 visible collars or 30 cm tall) stage of corn.

ULTIM® 75 DF HERBICIDE + DISTINCT* HERBICIDE

ULTIM® 75 DF Herbicide may be tank mixed with DISTINCT* Herbicide to control annual grasses, quackgrass, horsenettle, and many broadleaf weeds. Tank mix one water soluble bag/ha of ULTIM® 75 DF HERBICIDE and 285 grams of DISTINCT* Herbicide with a non-ionic surfactant at 0.25% v/v (2.5 L/1000 L) plus liquid urea ammonium nitrate (UAN) at 1.25% v/v (12.5 L/1000L) and apply postemergence to corn from the 2- to 6-leaf stage.

This tank mixture will control the weeds listed for control by ULTIM® 75 DF Herbicide alone, plus the weeds listed on the DISTINCT* label, plus horsenettle. Refer to the "ULTIM® 75 DF HERBICIDE USED ALONE" section of this label for more application instructions and restrictions on the use of ULTIM®75 DF. Refer to the DISTINCT* label for weeds controlled by DISTINCT* Herbicide and for additional application instructions and use precautions.

The tank mix of ULTIM® 75 DF Herbicide and DISTINCT* Herbicide should be applied to actively growing weeds at growth stages of emergence to 6-leaf (quackgrass at the 3- to 6-leaf stage, horsenettle at the 2- to 6-leaf). Do not apply after 6-leaf (4 visible collars or 30 cm tall) stage of corn.

ULTIM® 75 DF HERBICIDE + MARKSMAN* HERBICIDE

ULTIM® 75 DF Herbicide may be tank mixed with MARKSMAN* to control annual grasses, quackgrass and many broadleaf weeds. Tank mix one water-soluble bag/ha of ULTIM® 75 DF Herbicide and 2.5 L/ha of MARKSMAN* Herbicide with a non-ionic surfactant at 0.2% v/v (2 L/1000 L) and apply postemergence to corn up to the 5-leaf stage.

This tank mixture will control the weeds listed for control by ULTIM® 75 DF Herbicide alone (yellow foxtail suppression only), plus velvetleaf (most sensitive when small and actively growing), common ragweed and lamb's-quarters. Refer to the "ULTIM® 75 DF HERBICIDE USED ALONE" section of this label for more application instructions and restrictions on the use of ULTIM® 75 DF Herbicide. Refer to the MARKSMAN* Herbicide label for additional application instructions and use precautions.

The tank mix of ULTIM® 75 DF Herbicide and MARKSMAN* Herbicide must not be applied after the 5-leaf stage of corn.

ULTIM® 75 DF HERBICIDE + PARDNER* HERBICIDE

ULTIM® 75 DF Herbicide may be tank mixed with PARDNER* to control annual grasses, quackgrass and many broadleaf weeds. Tank mix one water soluble bag/ha of ULTIM® 75 DF Herbicide and 1.0 L/ha of PARDNER* Herbicide with a non-ionic surfactant at 0.2% v/v (2 L/1000 L) and apply postemergence to corn up to the 6-leaf stage (4 visible collars or 30 cm in height - leaf extended).

This tank mixture will control the weeds listed for control by ULTIM® 75 DF Herbicide alone, plus the weeds listed on the PARDNER* Herbicide label. Refer to the "ULTIM® 75 DF HERBICIDE USED ALONE" section of this label for more application instructions and restrictions on the use of ULTIM® 75 DF Herbicide. Refer to the PARDNER* label for weeds controlled by PARDNER* Herbicide and consult the label for additional application instructions and use precautions.

Applications of ULTIM® 75 DF Herbicide after the 6-leaf stage of corn (30 cm in height - leaf extended) must be delayed until after the 8-leaf stage of corn and before silking, when a height differential can be established between the corn and target weeds. When drop piping, ensure that the crop stand is UNIFORM so that all the spray solution is directed below the canopy to keep the spray out of the corn whorl.

ULTIM® 75 DF HERBICIDE + STRIKER* HERBICIDE

ULTIM® 75 DF Herbicide may be tank-mixed with STRIKER* Herbicide for added control of certain broadleaved weeds. Tank-mix one water-soluble bag/ha of ULTIM® 75 DF Herbicide and 280 grams/ha of STRIKER* with a non-ionic surfactant at 0.25% v/v (2.5 L/1000 L) in a total spray volume of 140-190 L/ha. Apply tank-mixture postemergence to corn up to the 6-leaf stage (4 visible collars or 30 cm in height - leaf extended). Do not add liquid fertilizer when tank-mixing ULTIM® 75 DF Herbicide with STRIKER* Herbicide.

This tank mix also offers suppression of field horsetail when used according to the labelled rates.

Observe relevant use directions, application instructions, precautions and high-risk environmental conditions listed on this label and the STRIKER* Herbicide label. The weeds controlled are those listed on this label in addition to those listed on the STRIKER* Herbicide label.

A 50 meter buffer zone appearing on the STRIKER* Herbicide label should be observed adjacent to aquatic habitats for this tank mix.

ULTIM® 75 DF HERBICIDE + STRIKER* HERBICIDE + AATREX* NINE-O

ULTIM® 75 DF Herbicide may be tank-mixed with STRIKER* Herbicide and AATREX* NINE-O for control of Eastern Black Nightshade and additional weeds listed on the ULTIM® 75 DF Herbicide and STRIKER* labels. Tank-mix one water-soluble bag of ULTIM® 75 and 280 grams/ha of STRIKER* Herbicide and 555 grams AATREX* NINE-O per hectare (500 gai/ha Atrazine) with a non-ionic surfactant at 0.25% v/v (2.5 L/1000 L) in a total spray volume of 140-190 L/ha. Apply tank-mixture postemergence to corn up to the 6-leaf stage (4 visible collars or 30 cm in height - leaf extended).

Observe relevant use directions, application instructions, precautions and high-risk environmental conditions listed on this label and the STRIKER* Herbicide and AATREX* NINE-O labels.

A 50 meter buffer zone appearing on the STRIKER* Herbicide label should be observed adjacent to aquatic habitats for this tank mix.

ULTIM® 75 DF HERBICIDE + CALLISTO* 480SC HERBICIDE

ULTIM® 75 DF Herbicide may be tank mixed with CALLISTO* 480SC Herbicide to control annual grasses, quackgrass and many broadleaf weeds. Tank mix one water-soluble bag/ha of ULTIM® 75 DF Herbicide and

0.21 L/ha of CALLISTO* 480SC Herbicide with the non-ionic surfactant, AGRAL* 90 at 0.2% v/v (2 L/1000 L), and apply postemergence to corn from the 3 to 6-leaf stage (4 visible collars or 30 cm in height - leaf extended).

This tank mixture will control the weeds listed for control by ULTIM® 75 DF Herbicide alone, plus the weeds listed on the CALLISTO* 480SC Herbicide label. Refer to the "ULTIM® 75 DF HERBICIDE USED ALONE" section of this label for more application instructions and restrictions on the use of ULTIM® 75 DF Herbicide. Refer to the CALLISTO* 480SC Herbicide label for weeds controlled by CALLISTO* 480SC Herbicide and consult the label for additional application instructions and use precautions.

The tank mix of ULTIM® 75 DF Herbicide and CALLISTO* 480SC Herbicide must not be applied after the 6-leaf stage (4 visible collars or 30 cm in height - leaf extended) of corn.

ULTIM® 75 DF HERBICIDE + CALLISTO* 480SC HERBICIDE + ATRAZINE

ULTIM® 75 DF Herbicide may be tank mixed with CALLISTO* 480SC Herbicide and Atrazine (either AATREX* NINE-0 or AATREX* Liquid 480) to control annual grasses, quackgrass and many broadleaf weeds. Tank mix one water-soluble bag/ha of ULTIM® 75 DF Herbicide, 0.21 L/ha of CALLISTO* 480SC Herbicide, and Atrazine (either 0.31 kg/ha of AATREX* Nine-0 or 0.58 L/ha of AATREX* Liquid 480) with the non-ionic surfactant, AGRAL* 90 at 0.2% v/v (2 L/1000 L), and apply postemergence to corn from the 3 to 6-leaf stage (4 visible collars or 30 cm in height - leaf extended).

This tank mixture will control the weeds listed for control by ULTIM® 75 DF Herbicide alone, plus the weeds listed on the CALLISTO* 480SC Herbicide label for the tank mix of CALLISTO* 480SC Herbicide + Atrazine (either AATREX Nine-0 or AATREX* Liquid 480). Refer to the "ULTIM® 75 DF HERBICIDE USED ALONE" section of this label for more application instructions and restrictions on the use of ULTIM® 75 DF Herbicide. Refer to the CALLISTO* 480SC Herbicide label for weeds controlled by the tank mix of CALLISTO* 480SC Herbicide + Atrazine (either AATREX Nine-0 or AATREX* Liquid 480) and consult the CALLISTO* 480SC Herbicide label and Atrazine (either AATREX Nine-0 or AATREX* Liquid 480) label for additional application instructions and use precautions.

The tank mix of ULTIM® 75 DF Herbicide, CALLISTO* 480SC Herbicide, and Atrazine (either AATREX Nine-0 or AATREX* Liquid 480) must not be applied after the 6-leaf stage (4 visible collars or 30 cm in height - leaf extended) of corn.

NOTE: Weeds that emerge after application of ULTIM® 75 DF Herbicide TANK MIXES will not be controlled. Application should be made when the majority of weeds have emerged.

MIXING INSTRUCTIONS:

Always add components to the spray tank in the following order: a) dry formulations in water soluble packaging, b) other dry materials and finally c) liquid materials.

1. Fill clean tank about 1/4 full with fresh water.
2. Turn on full agitation.
3. Add the proper amount of ULTIM® 75 DF Herbicide to the water in the spray tank with the agitator running. Maintain full agitation until water soluble bags are completely dissolved and product is fully dispersed. Continuous agitation is required to keep the product in suspension.
4. After ULTIM® 75 DF Herbicide has been well mixed, add the proper amount (if applicable) of AATREX* Nine-0, DISTINCT*, or STRIKER*, if applicable. Add slowly and agitate until completely dispersed.

5. While the mixture is in suspension add the proper amount (if applicable) of BANVEL* Herbicide, BANVEL* II Herbicide, MARKSMAN Herbicide, PARDNER* Herbicide, CALLISTO* 480 SC Herbicide, or AATREX* Liquid 480, to the spray tank with the agitator running. If Atrazine is being tank mixed with ULTIM® 75 DF Herbicide and STRIKER*, it should always follow the addition of STRIKER*.
6. After ULTIM® 75 DF Herbicide (and tank mixed herbicide if applicable) have been well mixed and is (are) in suspension add a recommended non-ionic surfactant such as CITOWETT* PLUS, AGRAL* 90, or AG-SURF* at 2 L per 1000 L spray solution (0.2% v/v) or SIDEKICK* at 2.5 L per 1000 L spray solution (0.25% v/v).
7. If the tank mix partner is DISTINCT* Herbicide add UAN at 12.5 L per 1000 L of water after the surfactant has been well mixed.
8. Fill the remainder of the spray tank.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be mixed.

Agitation is required for uniform mixing and application. The optimum water volume for ULTIM® 75 DF Herbicide application is 140-190 litres of water per hectare (minimum of 100 litres of water per hectare). Use a spray pressure of 175 - 275 kPa. Flat fan nozzles are recommended. Use 50 mesh filter screens or larger. Use spray preparation of ULTIM® 75 DF Herbicide within 24 hours or product degradation may occur resulting in a loss of weed control. Use vigorous agitation to thoroughly disperse spray mixtures that have been allowed to stand in the tank.

NOTE: ULTIM® 75 DF Herbicide will degrade in acidic or highly alkaline water. Mix no more than can be used in one day. If spraying is interrupted, thoroughly re-agitate the spray mixture before resuming spraying.

NOTE: Do not mix the soluble bags with liquid fertilizers, substances that contain boron or substances that release free chlorine. Mixing the water-soluble bags with any of these compounds will result in an insoluble substance in the tank.

MINOR USES

- **FOR USE ON FIELD CORN IN THE PROVINCE OF MANITOBA**
- **FOR USE ON LOWBUSH BLUEBERRIES IN EASTERN CANADA**

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than DuPont Canada and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. DuPont Canada itself makes no representation or warranty with respect to product performance (efficacy) and crop tolerance (phytotoxicity) of this product when used on the crops listed. Accordingly, the Buyer and User assume all liability arising, and agree to hold DuPont Canada harmless from any claims based on efficacy or phytotoxicity in connection with this use.

FOR USE ON FIELD CORN IN THE PROVINCE OF MANITOBA

For control of quackgrass, annual grasses and redroot pigweed, apply one water soluble bag of ULTIM® 75 DF Herbicide per hectare with a recommended non-ionic surfactant (CITOWETT* PLUS, AGRAL* 90

or AG-SURF*) at 2.0 litres per 1,000 litres of spray solution (0.2% v/v). (See "Mixing Instructions" section of the label).

Apply post-emergent to corn at the 1 to 4-leaf stage. Use only on the following corn hybrids:

PIONEER® Brand 39K72
PIONEER® Brand 39K73
PIONEER® Brand 39W54
PIONEER® Brand 39M27

Field corn, winter wheat and spring barley may be planted the year following application of ULTIM® 75 DF Herbicide. For all other crops, a field bioassay is recommended before planting. ULTIM® 75 DF Herbicide is more persistent in sandy soils, with low organic matter and pH 7 or higher. For these types of soils, a field bioassay is recommended before planting. Contact your local DuPont representative or the DuPont Hotline at 1-800-667-3925 for more information.

Apply with ground equipment only. Make only one application per growing season.

Please refer to the appropriate sections of the ULTIM® 75 DF Herbicide label booklet for additional use precautions, application and/or mixing instructions.

FOR USE ON LOWBUSH BLUEBERRIES IN EASTERN CANADA

For control of black bulrush, apply one water soluble bag of ULTIM® 75 DF Herbicide in 800 litres of water per 1 hectare with AGRAL* 90 at 2.0 litres per 1,000 litres of spray solution (0.2% v/v). See "Mixing Instructions" section of the label.

Apply ULTIM® 75 DF Herbicide to black bulrush in June of the prune year when the first flower heads are beginning to emerge from the bulrush tussock. ULTIM® 75 DF Herbicide should be applied as a directed spot spray to thoroughly wet the bulrush foliage. Weed control may be erratic or unsatisfactory from later applications and when the bulrush is under stress (e.g. drought stress).

Stunting and yield losses may occur when the blueberry plants are contacted by the spray.

Do not apply within 14 months of harvest.

Please refer to the appropriate sections of the ULTIM® 75 DF Herbicide label booklet for additional use precautions, application and/or mixing instructions.

SPRAYER CLEANUP:

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of ULTIM® 75 DF Herbicide as follows:

1. Drain tank; thoroughly hose down the interior surfaces of the tank; then flush tank, boom and hoses with clean water for a minimum of 5 minutes.
2. Fill the tank with clean water while adding one litre of household ammonia (containing a minimum of 3% ammonia) for every 100 litres of water. Flush the hoses, boom and nozzles with the cleaning solution and then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Again flush the hoses, boom and nozzles with the cleaning solution and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.

4. Repeat step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom.
6. PRIOR to using the sprayer for the next application, flush the tank, boom and hoses for 5 minutes with fresh water.

CAUTION:

DO NOT mix ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour that may cause eye, nose, throat and lung irritation. DO NOT clean equipment in an enclosed area.

USE OF INSECTICIDES:

For maximum crop safety, ULTIM® 75 DF Herbicide should be applied only to corn which has NOT been treated with a highly systemic organophosphorus soil insecticide, such as COUNTER*, THIMET*, or LORSBAN*. Application of ULTIM® 75 DF Herbicide to corn previously treated with FORCE* (pyrethroid) or FURADAN* (carbamate) will allow maximum crop safety.

DO NOT tank mix ULTIM® 75 DF Herbicide with any organophosphorus insecticide. DO NOT apply a foliar organophosphorus insecticide within 7 days before or after applying ULTIM® 75 DF Herbicide.

REPLANTING TO OTHER CROPS:

ULTIM® 75 DF Herbicide is degraded by natural soil processes, and field tests have shown that the following crops may safely be planted at the prescribed interval following corn that has been treated with ULTIM® 75 DF Herbicide.

Crop	Replanting Interval
Winter wheat	4 months
Spring barley, Canola, Soybeans, White beans, Red Clover, Sorghum, Field Corn	10 months

For other crops, a field bioassay is recommended before planting. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year.

FIELD BIOASSAY:

Select a representative area or areas of the field previously treated with ULTIM® 75 DF Herbicide to plant your bioassay crop(s). Be sure to consider factors such as size of field, soil texture, drainage and turn-around areas when selecting the site(s) that are most representative of the soil conditions in the field. On large fields, more than one site may be needed in order to obtain reliable results.

Plant the test strips perpendicular to the direction in which the field was sprayed. The strips should be long enough to cross the width of several spray swaths. Large test strip areas are more reliable than small ones.

Use standard tillage and seeding equipment to plant the bioassay. Prepare a seed bed and plant the crops and varieties you want the option of growing the following year. It is important to use the same planting time, conditions, techniques and cultural practices you normally use to plant and grow the bioassay crop(s). Also plant into an adjacent area not treated with ULTIM® 75 DF Herbicide to use as a comparison.

As the crop(s) emerges and grows, examine these key points in ULTIM® 75 DF Herbicide treated and non-treated areas:

crop stand
plant colour and vigour

root development
yield

rate of growth

Allow the bioassay crop(s) to grow to maturity while making your observations. Do not overspray the test strips with herbicides that may damage the bioassay crop(s). If the bioassay indicates that ULTIM® 75 DF Herbicide residues are still present, continue cropping only to those crops listed on the label and do not rotate to other crops until bioassay results indicate that susceptible crops are growing normally.

USE PRECAUTIONS:

Do not apply ULTIM® 75 DF Herbicide during periods of intense rainfall or to soils saturated with water. Do not apply directly to standing or running water. Do not apply in areas where surface water from the treatment site can run off to adjacent cropland, either planted or to be planted, or into streams, irrigation water or wells. Applications should only be made when there is no hazard of spray drift contaminating non-target land areas since very small quantities of the ULTIM® 75 DF Herbicide spray solution may severely injure susceptible crops during both growing and dormant periods.

As with any herbicide, overlaps or starting, stopping, slowing and turning while spraying may result in crop injury.

Crop injury may result if application is made to corn that has been stressed by abnormally hot, humid or cold weather conditions, frost, low fertility, drought, water saturated soil, compacted soil, previous pesticide applications, disease or insect damage.

Under extreme weather conditions, such as hot, dry weather, excessive moisture, or frost, weed control may be reduced.

Rainfall within 2-4 hours after application of ULTIM® 75 DF Herbicide may reduce weed control.

Because corn hybrids differ in their tolerance to herbicides, limit first use of ULTIM® 75 DF Herbicide to a small area of each hybrid prior to adoption as a field practice.

Overspray or drift to important wildlife habitats such as ponds, wetlands, streams, woodlots and shelterbelts should be avoided. Leave a 10-metre buffer zone between the last spray swath and bodies of water or wetlands. Leave a 5-metre buffer zone between the last spray swath and the terrestrial habitats listed above.

ULTIM® 75 DF Herbicide must not be applied within 30 days of harvest.

CAUTION:

Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, ULTIM® 75 DF Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to ULTIM® 75 DF Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the use of ULTIM® 75 DF Herbicide or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted.
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.

- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local DuPont representative or the DuPont hotline at 1-800-667-3925 for further information.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Keep water soluble bags in original outer bag or outer container away from moisture.

DISPOSAL:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For MYLAR® bag containers:

1. Make the empty bag unsuitable for further use.
2. Dispose of the bag in accordance with provincial requirements.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

The seller warrants that the purchase by the buyer and the use of this product, as such, will not infringe any Canadian patent.

REGISTERED TRADEMARKS:

®/tm Registered trademarks/trademarks of E.I. du Pont Company. E. I. du Pont Canada Company is a licensee. Member of CropLife.

* All other products mentioned are trademarks of their respective companies.

<<[Water-Soluble Bag]>>

ULTIM® 75 DF HERBICIDE

KEEP OUT OF REACH OF CHILDREN

AGRICULTURAL

SEE DIRECTIONS FOR USE BROCHURE FOR COMPLETE PRECAUTIONARY STATEMENTS

GUARANTEE: Rimsulfuron 37.5%
Nicosulfuron 37.5%

DRY FLOWABLE

NET CONTENTS: 33.7 GRAMS

WARNING: EYE IRRITANT

CAUTION: SKIN IRRITANT